

Dithyridus <sup>1</sup> (see Hemitryphis)—		591
<i>crustulum</i> , Bovallius, 1887,	.	269, 490, 591, 1464
<i>faba</i> , Dana, 1852,	.	591
<i>stellatus</i> , Bovallius, 1887,	.	591
<i>tenuimanus</i> , Bovallius, 1887,	.	591
Dodecas—		
<i>elongata</i> , Stebbing, 1883,	.	{ 461, 547, 1233, pls. cxxxix, cxl
Dryope—		
<i>crenatipluma</i> , Sp. Bate, 1862,	.	337, 343, 1145
<i>crenatipluma</i> , Bate and West-	.	343, 595
wood, 1863,	.	
<i>irrorata</i> , Sp. Bate, 1862,	.	337, 595, 1145
Dryopoides—		
<i>westwoodi</i> , Stebbing, 1888,	.	1146, pl. cxxii
Dulichia—		
<i>curticauda</i> , Boeck, 1870,	.	396
<i>falcata</i> , Sp. Bate, 1857,	.	293, 306, 396, 494
<i>hirticornis</i> , G. O. Sars, 1876,	.	459, 571
<i>macera</i> , G. O. Sars, 1879,	.	498, 571
<i>Malmgreni</i> , Iarzynsky, 1870,	.	403
<i>monacantha</i> , Metzger, 1875,	.	445, 542
<i>Nordlandica</i> , Boeck, 1870,	.	396
<i>porrecta</i> , Sp. Bate, 1857,	.	{ 293, 306, 396, 445, 494 542
<i>septentrionalis</i> , G. O. Sars,	.	498, 571
1879,	.	
<i>spec. dubia</i> , Metzger, 1875,	.	445
<i>spinossissima</i> , Krøyer, 1845,	.	{ 213, 217, 302, 358, 396 1634
<i>tuberculata</i> , Boeck, 1870,	.	396, 498, 571
Dyopodus (see Dulichia)—		
<i>falcatus</i> , Sp. Bate, 1857,	.	293
<i>porrectus</i> , Sp. Bate, 1857,	.	293
Egidia <sup>2</sup> (see Urothoe)—		
<i>pulchella</i> , A. Costa, 1853,	.	274, 297
Eiscladus (see Photis)—		
<i>longicaudatus</i> , Bate and West-	.	341, 396, 430, 466
wood, 1862,	.	
Elamis (see Amphithoe)—		
<i>viridis</i> , Leach, MS., in White,	.	222
1847,	.	
Elasmocerus—		
<i>Speciosus</i> , Costa, 1851,	.	248, 250
Elasmopus—		
<i>brevicaudata</i> , Boeck, 1872,	.	1018
<i>delaplata</i> , Stebbing, 1888,	.	1025, pl. xcix
<i>erythræus</i> , Stebbing, 1888,	.	516
<i>latipes</i> , Boeck, 1870,	.	395, 594, 595, 1649
<i>rapax</i> , A. Costa, 1853,	.	274, 297, 469, 560, 1649
<i>subcarinata</i> , Stebbing, 1888,	.	1019, 1637, pl. xcvi
Enone—		
<i>punctata</i> , Risso, 1826,	.	129, 249, 560
Ephippiphora <sup>3</sup> —		
<i>Kroyeri</i> , White, 1847,	.	224, 500, 555, 587
Epidesura—		
<i>compressa</i> , Boeck, 1860,	.	322
Epimeria—		
<i>conspicua</i> , Stebbing, 1883,	.	547, 878
<i>cornigera</i> , Bate and Westwood,	.	{ 45, 223, 243, 250, 292 1868,
<i>loricata</i> , G. O. Sars, 1879,	.	{ 297, 305, 318, 322, 369 395, 417, 569, 1648
<i>sp.</i> , Catta (Chatin), 1878,	.	{ 498, 531, 547, 569 878, pl. lxviii
<i>tricristata</i> , <sup>4</sup> A. Costa, 1851,	.	475
Ericthonius (often spelt Erichthonius)—		
<i>abditus</i> , Chevreux, 1887,	.	595
<i>bidens</i> , A. Costa, 1853,	.	274, 299, 367, 560, 595
<i>difformis</i> , M.-Edw., 1830,	.	{ 142, 188, 200, 277, 284 292, 299, 312, 336, 358 396, 437, 546, 595
<i>longimanus</i> , S. I. Smith,	.	396, 1132
<i>? macrodactylus</i> , Dana, 1852,	.	255, 268
<i>megalops</i> , G. O. Sars, 1885,	.	498, 571
<i>? minax</i> , 1888,	.	437
<i>pugnax</i> , Dana, 1852,	.	255, 267
<i>rapax</i> , Stimpson, 1857,	.	303, 410
Eriopis—		
<i>elongata</i> , Bruzelius, 1859,	.	313, 395
Erpetoramus—		
<i>Costae</i> , De Natale, 1850,	.	248, 347, 1623
Erythrocephalus—		
<i>cæcus</i> , Tilesius, 1819,	.	109
<i>macrophthalmus</i> , de Quatrefages,	.	275
1853,	.	
<i>melanophthalmus</i> , Tilesius, 1819,	.	109, 167
Euonyx—		
<i>chelatus</i> , Norman, 1867,	.	370, 673
<i>normani</i> , Stebbing, 1888,	.	669, pl. xix
Eupheus—		
<i>ligioides</i> , Risso, 1816,	.	97, 105, 129, 138
Eupronoë—		
<i>armata</i> , Claus, 1879,	.	241, 269, 492, 591, 1519
<i>atlantica</i> , Stebbing, 1888,	.	1519
<i>brunnea</i> , Bovallius, 1887,	.	269, 492, 591, 1510
<i>inscripta</i> , Stebbing, 1888,	.	1510, pl. clxxxvii
<i>intermedia</i> , Stebbing, 1888,	.	1517, pl. clxxxviii
<i>macrocephala</i> , Bovallius, 1887,	.	591
<i>maculata</i> , Claus, 1879,	.	241, 492, 591, 1510
<i>minuta</i> , Claus, 1879,	.	492, 591, 1516
<i>ornata</i> , Bovallius, 1887,	.	591
<i>pacifica</i> , Stebbing, 1888,	.	1513
<i>serrata</i> , Claus, 1887,	.	492
Eurymera—		
<i>monticulosa</i> , Pfeffer, 1888,	.	1653

<sup>1</sup> The suggestion, p. 97, that "*Typhis ovoides*, Riss," should be called *Dithyridia ovoides*, as well as the suggestion, p. 500, that *Platyscelus intermedius*, Thomson, should be called *Dithyridia intermedius*, is withdrawn for the reasons mentioned on pp. 1463, 1464.

<sup>2</sup> Boeck, De Skand. og. Arkt. Amph., p. 58, spells this name *Aegidia*.

<sup>3</sup> Since *Ephippiphora* is preoccupied, the suggestion made on p. 177 that it will take precedence of Boeck's *Socarnes* must be cancelled, and with it the name "*Ephippiphora vahlii*, Krøyer," falls to the ground.

<sup>4</sup> Bate and Westwood, Brit. Ses. Crust., vol. ii. p. 528, identify *Acanthonotus owenii*, Sp. Bate, and *Epimeria tricornisata*, Costa, with *Epimeria cornigera* (Fabricius), therefore anticipating Boeck, who independently took the same view in 1870.