

| | |
|---|------------------------------|
| Gammarus— <i>continued</i> . | |
| <i>tenuicornis</i> , Stimpson, 1855-6, | 288 |
| <i>tenuimanus</i> , Sp. Bate, 1862, | 335 |
| <i>tenuis</i> , Dana, 1852, | 255 |
| <i>testaceus</i> , Dybowski, 1874, | 427 |
| <i>thaumops</i> , Grimm, 1880, | 509 |
| <i>Torelli</i> , Goës, 1865, | 357, 368, 395 |
| <i>toxophthalmus</i> , Dybowski, 1874, | 428 |
| <i>truncatus</i> , Viviani, 1805, | 76 |
| <i>tuberculatus</i> , Dybowski, 1874, | 428 |
| <i>tunetanus</i> , E. Simon, 1885, | 573 |
| <i>unguiserratus</i> , A. Costa, 1853, | 274, 298 |
| { <i>Ussolzevii</i> , Dybowski, 1874, | 428 |
| { <i>Ussolzevii</i> , Dybowski, 1874, | 428 |
| <i>var. abyssorum</i> , Dybowski, 1874, | 428 |
| <i>validus</i> , Dana, 1852, | 255, 267 |
| <i>Veneris</i> , Heller, 1865, | 359, 388 |
| <i>verrucosus</i> , Gerstfeldt, 1858, | 309, 427 |
| <i>Victorii</i> (see <i>Godlewskii</i> , Dybowski, 1874), | 428 |
| <i>violaceus</i> (see <i>fasciatus</i> , Costa), | 206 |
| <i>violaceus</i> , Dybowski, 1874, | 428 |
| <i>var. virescens</i> , Dybowski, 1874, | |
| <i>virescens</i> (see <i>violaceus</i> , Dybowski, 1874), | |
| <i>viridis</i> , Dybowski, 1874, | 428 |
| <i>var. canus</i> , Dybowski, 1874, | 428 |
| <i>var. olivaceus</i> , Dybowski, 1874, | 428 |
| <i>vittatus</i> , Dybowski, 1874, | 428 |
| <i>vortex</i> , Dybowski, 1874, | 428 |
| <i>Wagii</i> , Dybowski, 1874, | 428 |
| <i>WahlII</i> , Dybowski, 1874, | 428 |
| <i>var. platycercus</i> , Dybowski, 1874, | 428 |
| <i>zebra</i> , Rathke, 1843, | 204, 252, 285, 312 |
| <i>zebra</i> , Dybowski, 1874, | 428 |
| <i>Zete</i> , White, 1847, | 223 |
| <i>ZienkowiezII</i> , Dybowski, 1874, | 428 |
| Gitana— | |
| <i>Sarsi</i> , Boeck, 1870, | 394, 1650 |
| <i>rostrata</i> , Boeck, 1870, | 394 |
| <i>Glauconome</i> (see <i>Unciola</i>)— | |
| <i>Krøyeri</i> , Boeck, 1870, | 370, 374, 396, 1170 |
| <i>leucopis</i> , Krøyer, 1845, | { 212, 213, 217, 302 |
| <i>leucopis</i> , Krøyer, 1845, | { 319, 358, 374, 396 |
| <i>leucopis</i> , Krøyer, 1845, | { 424, 546, 1634 |
| <i>petalocera</i> , G. O. Sars, 1879, | 459, 498, 571 |
| <i>planipes</i> , G. O. Sars, 1876, | 370, 459, 498 |
| <i>Steenstrupi</i> , Boeck, 1870, | 396, 1170, ¹ 1650 |
| Glosscephalus— | |
| <i>Milne-Edwardsi</i> , Bovallius, 1887, | 590 |
| <i>spiniger</i> , Bovallius, 1887, | 590 |
| <i>Glycera</i> — | |
| <i>tenuicornis</i> , Haswell, 1880, | 511 |
| <i>Glycerina</i> — | |
| <i>affinis</i> , Chilton, 1884, | 551 |
| <i>tenuicornis</i> , Haswell, 1882, | 512, 551 |
| Goësia— | |
| <i>depressa</i> , Boeck, 1870, | 358, 396 |
| Goplana— | |
| <i>ambulans</i> , Wrzeźniowski, 1879, | 217, 502 |

Goplana—*continued*.

| | |
|---|--------------------------------------|
| <i>polonica</i> , Wrzeźniowski, 1879, | { 217, 501, 502, 503-507 |
| <i>polonica</i> , Wrzeźniowski, 1879, | { 519, 532 |
| Gossea— | |
| { <i>microdentopa</i> , Sp. Bate, 1862, | 334 |
| { <i>microdentopa</i> , Sp. Bate, 1862, | 334, 504, 543 |
| <i>Grayia</i> — | |
| <i>imbricata</i> , Sp. Bate, 1862, | 266, 332, 425, 435, 498 |
| <i>Pugettensis</i> , Sp. Bate, 1862, | 266, 332 |
| Grubia— | |
| <i>taurica</i> , Czerniawski, 1868, | 377 |
| <i>Guerinia</i> (see <i>Trischizostoma</i>)— | |
| <i>nicæensis</i> , Costa, 1853, | 272, 369 |
| Guerneæ— | |
| <i>coalita</i> , Chevreux, 1887, | 386, 595 |
| <i>lævis</i> , Chevreux, 1887, | 596 |
| Halice— | |
| <i>abyssi</i> , Boeck, 1870, | 394, 542 |
| <i>grandicornis</i> , Boeck, 1870, | 394, 542 |
| Haliceion— | |
| ? <i>latipes</i> , G. O. Sars, 1882, | 540, 599 |
| <i>longicaudatus</i> , Boeck, 1870, | 395, 540 |
| Halimedon— | |
| <i>brevicalcar</i> , Boeck, 1870, | 356, 395 |
| <i>longimanus</i> , Boeck, 1870, | 394 |
| <i>megalops</i> , G. O. Sars, 1882, | 540 |
| { <i>Mølleri</i> , Boeck, 1870, | 394 |
| { <i>Mülleri</i> , Boeck, 1876, | 394 |
| <i>obtusifrons</i> , Hansen, 1887, | 1644 |
| <i>Saussurei</i> , Boeck, 1870, | 394, 546 |
| <i>schneideri</i> , Stebbing, 1888, | 839, pl. lix |
| Halirages— | |
| <i>bispinosus</i> , Boeck, 1870, | 292, 314, ² 319, 395, 548 |
| <i>borealis</i> , Boeck, 1870, | 395 |
| <i>fulvocinctus</i> , Boeck, 1870, | { 319, 345, 356, 395, 468 |
| <i>fulvocinctus</i> , Boeck, 1870, | { 546, 569, 600, 901 |
| <i>fulvocinctus</i> , Boeck, 1870, | { 1634 |
| <i>huxleyanus</i> , Stebbing, 1888, | 334, 902, pl. lxxiii |
| <i>inermis</i> , G. O. Sars, 1882, | 540 |
| <i>maculatus</i> , Stuxberg, 1880, | 523 |
| <i>megalops</i> , G. O. Sars, 1882, | 540 |
| <i>quadridentatus</i> , G. O. Sars, 1876, | 458, 569 |
| <i>tridentatus</i> , Boeck, 1870, | { 314, 356, 395, 458, 540 |
| <i>tridentatus</i> , Boeck, 1870, | { 932 |
| Haplocheira— | |
| <i>barbimanus</i> , Stebbing, 1888, | 500, 511, 551, 565, 1177 |
| <i>plumosa</i> , Stebbing, 1888, | 1172, pl. cxxvi |
| <i>typica</i> , Haswell, 1880, | 500, 511, 565 |
| Haploops— | |
| <i>carinata</i> , Liljeborg, 1855, | 270, 285, 314, 395 |
| <i>lævis</i> , Hoek, 1882, | 534, 600 |
| <i>lineata</i> , Stuxberg, 1880, | 523 |
| <i>setosa</i> , Boeck, 1870, | 395, 531 |
| <i>tubicola</i> , Liljeborg, 1855, | { 270, 285, 314, 357, 373 |
| <i>tubicola</i> , Liljeborg, 1855, | { 385, 395, 525, 600, 1634 |
| <i>Harmonia</i> — | |
| <i>crassipes</i> , Haswell, 1880, | 512, 565 |
| <i>Harmophia</i> — | |
| <i>Krøyeri</i> (Boeck, MS.), 1877, | 466 |

¹ On p. 1170, line 17, for *steenstrupii* read *steenstrupi*.² On p. 314, line 18, for *bispinosa* read *bispinosus*.