

<i>Leptocheirus</i> — <i>continued</i> .	
<i>pinguis</i> , Stebbing, 1888,	279
<i>trieristatus</i> , Stebbing, 1888,	595
<i>Leptochela</i> —	
<i>leptochela</i> , Boeck, 1876,	373, 453
<i>Leptocotis</i> —	
<i>ambobus</i> , Stebbing, 1888,	{ 1594, 1597, fig. 31, pl. ccv
<i>Lindströmi</i> , Bovallius, 1887,	590, 1596
<i>mindanaonis</i> , Stebbing, 1888,	1598, pl. cciv, c
<i>spinifera</i> , Streets, 1877,	{ 470, 493, 590, 1593 1597
<i>tenuirostris</i> , Bovallius, 1887,	590, 1593, 1596
<i>Leptomera</i> (see Proto)—	
<i>pedata</i> , Lamarck, 1818,	{ 105, 136, 138, 163, 175 191, 203, 205, 223, 281 284, 319, 327, 342, 421 457, 1621
<i>rubra</i> , Lamarck, 1818,	105, 175, 203
<i>ventricosa</i> , Desmarest, 1823-5,	{ 119, 123, 138, 183, 191 192, 203
<i>Leptothoe</i> (see Mæra)—	
<i>Danæ</i> , Stimpson, 1854,	277
<i>Lestrigonus</i> (see Hyperia)—	
<i>bengalensis</i> , Giles, 1887,	1391, 1643
<i>Boeckii</i> , Boeck, 1860,	143, 321
<i>exulans</i> , Krøyer, 1838,	{ 179, 189, 302, 321, 354 358, 382, 494, 1645
{ <i>Fabrei</i> , M.-Edw., 1830,	143, 175, 184, 189
{ <i>Fabreii</i> , M.-Edw., 1840,	{ 143, 189, 268, 290, 354 1390
<i>Fabricii</i> ?, Kinahan, 1859,	317, 354, 426
<i>falcatus</i> , Kinahan, 1863,	344
<i>ferus</i> , Dana, 1852,	268, 588
<i>fuscus</i> , Dana, 1852,	268, 589, 1403
<i>Gaudichaudii</i> , Sp. Bate, 1862,	189
<i>Kinahani</i> , Sp. Bate, 1862,	{ 337, 354, 358, 382, 421 430, 465, 494
<i>mediterraneus</i> , Costa, 1865,	354, 561, 588
<i>rubescens</i> , Dana, 1852,	268, 354, 469, 588
<i>spinidorsalis</i> , Sp. Bate, 1877,	473
<i>Leucothoe</i> ,	
<i>affinis</i> , Stimpson, 1855-6,	84, 771
<i>antarctica</i> , Pfeffer, 1888,	288
<i>articulata</i> (see <i>articulosa</i>),	1653
<i>articulosa</i> , Leach, 1813-14,	{ 106, 120 84, 90, 122, 183, 188 223, 249, 278, 283, 284 315, 367, 442, 545
<i>brevidigitata</i> , Miers, 1884,	555, 782
<i>clypeata</i> , Krøyer, 1842,	{ 200, 214, 217, 236, 284 252, 302, 315, 373, 394 446
<i>clypeata</i> ?, Bruzelius, 1859,	314, 356
<i>commensalis</i> , Haswell, 1880,	511, 555, 565, 776
<i>crassimana</i> , Kossmann, 1880,	516, 555
<i>denticulata</i> , Costa, 1853,	{ 183, 274, 299, 329, 367 442, 545
<i>diemenensis</i> , Haswell, 1880,	511, 556, 565

Leucothoe—*continued*.

<i>hindersi</i> , Stebbing, 1888,	779, pl. xlvi
<i>furina</i> , Audouin, 1825,	{ 93, 120, 127, 136, 163 176, 284, 292, 295, 305 388, 542, 560
<i>glacialis</i> , Krøyer, 1842,	200, 214, 217, 284, 394
<i>gracilis</i> , Haswell, 1880,	511, 556, 565
<i>grandimanus</i> , Stimpson, 1854,	278
<i>Lilljeborgii</i> , Boeck, 1860,	322
<i>miersi</i> , Stebbing, 1888,	772, pl. xlvi
<i>norvegica</i> , Liljeborg, 1850,	{ 236, 251, 284, 315, 319 373
<i>novæ-hollandiæ</i> , Haswell, 1880,	512, 555
<i>Parthenopæa</i> , Costa, 1851,	249, 299
<i>phyllonyx</i> , M. Sars, 1859,	314, 319, 395
<i>procera</i> , Sp. Bate, 1857,	292, 295, 305
<i>Richiardi</i> (Lesson, MS.), Carus, }	560
1885,	
<i>spinicarpa</i> , Boeck, 1860,	{ 75, 84, 299, 322, 367 394, 516, 542, 555, 565
var. <i>commensalis</i> , Haswell, 1885,	565
var. <i>diemenensis</i> , Haswell, 1885,	565
var. <i>gracilis</i> , Haswell, 1885,	565
<i>stylifera</i> , Stimpson, 1855-6,	288
<i>traillii</i> , G. M. Thomson, 1882,	1637
<i>tridens</i> , Stebbing, 1888,	777, pl. xlvii
<i>Liljeborgia</i> (often spelt <i>Lilljeborgia</i>)—	
<i>æquabilis</i> , Stebbing, 1888,	988
<i>æquicornis</i> , G. O. Sars, 1876,	458, 570
<i>consanguinea</i> , Stebbing, 1888,	980, pl. xci
<i>fissicornis</i> , Boeck, 1870,	319, 324, 357, 394
<i>haswelli</i> , Stebbing, 1888,	565, 985, pl. xcii
<i>Kinahani</i> , ¹ Boeck, 1876,	333
<i>Normanni</i> , Stebbing, 1874,	204, 434, 496
<i>pallida</i> , Sp. Bate, 1862,	{ 292, 313, 322, 333, 357 394, 442, 545, 984
<i>shetlandica</i> , ² Bate and Westwood, }	204, 340, 434, 496, 1628
1862,	
<i>Limnoria</i> (see <i>Chelura</i>)—	
<i>xylophaga</i> , Hesse, 1868,	383
<i>Liparis</i> —	
<i>lobata</i> , Bosc, 1802,	1618
<i>Liriope</i> —	
<i>pygmæa</i> , Rathke, 1843,	204
<i>Lonchomerus</i> —	
<i>gracilis</i> , Sp. Bate, 1857,	292
<i>Lusya</i> (? <i>Erichthonius</i>)—	
<i>algensis</i> , Nardo, 1847,	220, 390, 1157
<i>Lycæa</i> —	
<i>longicornuta</i> , Stebbing, 1888,	1566, 1643
{ <i>malinii</i> , G. O. Sars, 1882,	393, 431, 541, 593
{ <i>nasuta</i> , Claus, 1879,	492, 590
<i>ochracea</i> , Dana, 1852,	269, 431, 590
<i>pauli</i> , Stebbing, 1888,	1566
<i>pulex</i> , Marion, 1874,	431, 492, 541, 561, 590, 1567
<i>robusta</i> , Claus, 1879,	431, 492, 561, 590, 1567
<i>serrata</i> , Claus, 1879,	492, 590
<i>similis</i> , Claus, 1879,	492, 590

¹ See also Chevreux, *Nouvelles espèces de Crustacés Amphipodes du sud-ouest de la Bretagne*. Assoc. Franç. pour l'avancement des sciences, Congrès de Toulouse, 1887 (1880).

² On p. 496, line 25, for Sp. Bate, read Bate and Westwood.