

Parelasomopus—			
<i>suluensis</i> , Stebbing, 1888,	1029, pl. c		
Pariambus (see <i>Podalirius</i>)—			
<i>Krøyeri</i> , 1888, }			
<i>minutus</i> , 1888, }		1268	
<i>typicus</i> , 1888, }			
<i>Pediculus</i> —			
<i>ceci</i> , Seba, 1734,		7, 11, 201	
<i>Peltocoxa</i> —			
<i>Marioni</i> , Catta, 1875,			441
<i>Pepredo</i> —			
<i>potamogeti</i> , Rafinesque, 1817,			100
<i>Pereionotus</i> —			
<i>testudo</i> , Bate and Westwood, 1862,	81, 341, 441, 1638		
<i>Phædra</i> —			
<i>antiqua</i> , Sp. Bate, 1859,			311
<i>Kinahani</i> , Sp. Bate, 1862,	333, 435, 1725		
<i>spinifera</i> , Sp. Bate, 1862,			277, 333
<i>Phasmatocarcinus</i> —			
<i>discophthalmus</i> , Tilesius, 1819,		109, 135	
<i>glaucus</i> , Tilesius, 1819,		109, 135	
<i>Pherusa</i> —			
<i>australis</i> , Haswell, 1880,			514
<i>Barretti</i> , Sp. Bate, 1862,			334
<i>bicuspis</i> , ¹ Sp. Bate, 1862,	179, 314, 1635		
<i>bispinosa</i> , Nebeski, 1880,			520
<i>cærulea</i> , G. M. Thomson, 1886,		586, 1648	
<i>cirrus</i> , ¹ Sp. Bate, 1862,		334, 1635	
<i>costata</i> , Sp. Bate, 1862,			141
<i>elegans</i> , ² Sp. Bate, 1862,			314
<i>fucicola</i> , Leach, 1814,	{ 86, 90, 91, 123, 131 141, 176, 187, 221, 222 298, 365, 547, 578, 1625		
<i>inermis</i> , Czerniavski, 1868,			379
<i>lævis</i> , Haswell, 1880,			511
<i>neo-zelanica</i> , Thomson and Chil- } <i>ton</i> , 1886,			586
<i>novæ-zealandiæ</i> , G. M. Thom- } <i>son</i> , 1879,		499, 586	
<i>podura</i> , Sp. Bate, 1862,			56
<i>pontica</i> , Czerniavski, 1868,			379
<i>pulchella</i> , Sp. Bate, 1862,			314
<i>tricuspis</i> , Stimpson, 1863,			345, 356
<i>Phlias</i> —			
<i>rissoanus</i> , Sp. Bate, 1862,	81, 332, 341, 441		
<i>serratus</i> , Guérin, 1836,	{ 81, 165, 176, 186, 341 1637		
<i>Phorcorrhaphis</i> —			
<i>edwardsi</i> , Stebbing, 1888,		1455, pl. clxxxi	
<i>zamboangæ</i> , Stebbing, 1888,		1452, pl. clxxx	
<i>Phorcus</i> (now <i>Phorcorrhaphis</i>)—			
<i>hyalocephalus</i> , Dana, 1852,		269, 589, 1457	
<i>Lovéni</i> , Bovallius, 1887,			589, 1457
<i>Raynaudii</i> , M.-Edw., 1840,		142, 189, 269	
<i>Reynaudii</i> , M.-Edw., 1830,		142, 175, 184, 589, 1457	
<i>Photis</i> —			
<i>brevicaudata</i> , Stebbing, 1888,			1068, pl. cviii
<i>longicaudata</i> , Meinert, 1877,			466
<i>Lütkeni</i> , Boeck, 1870,			396
<i>macrocarpus</i> , Stebbing, 1888,			1064, pl. cvii
<i>Reinhardi</i> , Krøyer, 1842,	{ 199, 271, 301, 313, 341 358, 396		
<i>Reinhardti</i> , Lütken, 1875,			1634
<i>tenuicornis</i> , G. O. Sars, 1882,			540
<i>Phoxocephalus</i> —			
<i>bassi</i> , Stebbing, 1888,		811, 1637, pl. liv	
<i>kergueleni</i> , Stebbing, 1888,			816, pl. lv
<i>Phoxus</i> (now <i>Phoxocephalus</i>)—			
<i>Batei</i> , Haswell, 1880,		511, 815, 1637	
<i>erythrophthalmus</i> , Catta, 1875,			441, 561
<i>falcatus</i> , G. O. Sars, 1882,			394, 538
<i>fusiformis</i> , Stimpson, 1854,			279, 431
<i>geniculatus</i> , Stimpson, 1855-6,			287
<i>grandis</i> , Stimpson, 1857,			303
<i>Holbölli</i> , Krøyer, 1842,	{ 198, 214, 292, 357, 394 465, 568, 1634		
<i>Holbolli</i> , Sp. Bate, 1857,			292, 295
<i>Holbøllii</i> , Reinhardt, 1857,			301, 431
<i>Kroyeri</i> , Stimpson, 1854,			279, 431
<i>Kroyerii</i> , Sp. Bate, 1857,			279, 292, 295
<i>maculatus</i> , Chevreux, 1888,			1650
<i>obtusus</i> , Stimpson, 1855-6,			287
<i>oculatus</i> , G. O. Sars, 1879,		498, 568, 1650	
<i>plumosus</i> , Krøyer, 1842,	{ 198, 214, 279, 301, 357 394, 431, 819, 1638		
<i>rostratus</i> , Boeck, 1876,			266
<i>simplex</i> , Sp. Bate, 1857,			279, 292, 295
<i>simplex</i> , Boeck, 1870,			394, 538
<i>villosus</i> , Haswell, 1880,			511, 815
<i>Phreatoicus</i> —			
<i>typicus</i> , Chilton, 1883,			543, 587
<i>Phronima</i> —			
<i>atlantica</i> , Guérin, 1836,	{ 162, 165, 175, 190, 223 269, 281, 339, 413, 437 469, 542, 589, 1351 1624, pl. clx.		
<i>atlantica</i> , White, 1847, ³			223, 337
<i>Borneensis</i> , Sp. Bate, 1862,			337, 413, 469
<i>bucephala</i> , Giles, 1887,			1642
<i>Cocconi</i> , de Natale, 1850,			1624
<i>Colletti</i> , Bovallius, 1887,			589
<i>custos</i> , Risso, 1816,	{ 97, 106, 122, 124, 135 137, 183, 190, 248, 272 339, 469		
<i>clongata</i> , Claus, 1862,			339, 405, 470, 1361
<i>megalodous</i> , Stebbing, 1888,			1353, pl. clxii, A
<i>neo-zelanica</i> , Thomson and Chil- } <i>ton</i> , 1886,			587
<i>novæ-zealandiæ</i> (?), Powell, 1875, } pl. clxi, B	{ 448, 587, 589, 1356		

¹ This is not Krøyer's *Amphithoe bicuspis*; that view, though accepted at p. 179, is corrected on p. 1635. Bate and Westwood give *Pherusa cirrus* as a synonym of their *Pherusa bicuspis*.

² Brit. Mus. Catal. Amph. Crust., p. 377, for *Paramphithoe elegans*, Bruzelius (see p. 314).

³ The Brit. Mus. Catal. Amph. Crust. refers to White's Catalogue of 1850 by mistake for that of 1847.