

Protella—continued.

Danae, Kossmann, 1880,	515
dentata, Haswell, 1885,	535
{ echinata, Mayer, 1882,	535
{ echinimana, ¹ Mayer, 1882,	536
gracilis, Dana, 1852,	265, 535, 565, 1245
Haswelliana, Mayer, 1882,	535, 565, 1226
intermedia, Czerniavski, 1868,	375
longispina, Sp. Bate, 1859,	290, 306, 322
major, Haller, 1879,	478
Phasma, Sp. Bate, 1862,	{ 74, 124, 151, 195, 212 322, 397, 421, 478, 479 515, 521, 527, 535, 578 1244
subspinosa, Kossmann, 1880,	515
typica, Czerniavski, 1868,	375
Protellopsis—	
kerueleni, Stebbing, 1888,	1241, pl. cxlii
Proto (see Phtisica), ² —	
brunneovittata, Haller, 1879,	749, 535
condylata, Haswell, 1885,	564
? cornigera, Mayer, 1882,	535, 565
{ elongata, Sp. Bate, 1862,	265
{ elongatus, Dana, 1852,	265
{ Goodsiri, Stebbing, 1876,	459, 479
{ Goodsirii, Sp. Bate, 1857,	293, 306, 396, 459, 527
Novae-Hollandiae, Haswell, 1880,	511, 535, 566, 1230
{ Pedata, Leach, 1814,	{ 86, 91, 243, 322, 459 479, 527
{ pedatum, W. Thompson, 1844,	208, 223
{ pedatus, Fleming, 1823,	119, 157, 254
spinosa, Haswell, 1885,	564
ventricosa, Boeck, 1870,	{ 19, 83, 105, 158, 191 223, 293, 322, 327, 396 459, 479, 483, 525, 535 548, 596
Protomedeia—	
fasciata, Krøyer, 1842,	{ 199, 286, 301, 313, 357 396, 466, 561, 1634
fasciata, Costa, 1864,	346, 561
fimbriata, ³ Sp. Bate, 1862,	
guttata, Grube, 1866,	366, 561, 1627
{ hirsutimana, Bate and West- wood, 1862,	545, 561, 1628
{ var. massiliensis, Catta, 1885,	561
hirsutimanus, Sp. Bate, 1862,	334, 346, 366, 466
longimana, Boeck, 1870,	396
massiliensis (see hirsutimana),	561
Nordmannii, Sp. Bate, 1862,	186, 200
pectinata, Norman, 1869,	1628
pilosa, Sp. Bate, 1862,	{ 366, 485, 486, 561, 1627 1639
pinguis, Sp. Bate, 1862,	279
Whitei, Sp. Bate, 1862,	334, 1628
Proton—	
pedatum, Desmarest, 1823-5,	123, 149

Psammylla—

littoralis, Rafinesque, 1817,	100
Pseudolycea—	
pachypoda, Claus, 1879,	493, 590
Pseudophthalmus (see Ampelisca)—	
ingens, Stimpson, MS., 1862,	279, 332
limicola, ⁴ Stimpson, 1854,	279
pelagicus, Stimpson, 1854,	279
Pterygocera (see Haustorius)—	
arenaria, Latreille, 1829,	{ 39, 138, 163, 444, 457 474
Pterygocerus—	
arenarius, Latreille, 1825,	126
Ptilocheirus (see Leptocheirus)—	
hirsutimanus, Norman, 1869,	595, 1628
pectinatus, Chevreux, 1887,	595
Pilosus, Meinert, 1877,	466
pinguis, Stimpson, 1854,	279, 432, 436, 437
tricristatus, Chevreux, 1887,	594, 595
Pulex—	
cancriformis, Strøm, 1762,	24
cornutus, Ray, 1710,	9
fluviatilis, Ray, 1710,	10
fluviatilis, Linnæus, 1751,	14
marinus, Rondelet, 1554,	3, fig. 1, 4, 5, 9, 13
Pycnogonum—	
ceti, J. C. Fabricius, 1798,	65, 129, 1618
Pyctilus—	
brasiliensis, Dana, 1852,	268
macrodaetylus, Dana, 1852,	255, 268, 375, 560
pugnax, Dana, 1852,	255, 268, 375, 560
Reptorramphus (see Erpetorramphus)—	
Costæ, Boeck, 1872,	1624
Rhabdonectes (see Rhabdosoma)—	
armatus, Bovallius, 1887,	591
Whitei, Bovallius, 1887,	591
Rhabdosoma—	
armatum, Adams and White,	{ 191, 224, 225, 308, 338 406, 485, 493, 591 1607, 1610, fig. 32 1643
batei, Kingsley, 1884,	554
brevicaudatum, Stebbing, 1888,	1607, 1612, pl. ccviii
curvicorne, Sp. Bate, 1862,	1608
investigatoris, Giles, 1887,	1609, 1643
longirostris, Streets, 1878,	485, 1608
Whitei, Sp. Bate, 1862,	{ 338, 406, 485, 493, 591 1609, 1611
Rhachotropis (see Tritropis)—	
aculeata, S. I. Smith, 1883,	{ 49, 113, 161, 216, 394 423, 546
aculeatus, Stebbing, 1888,	954
avirostris, Stebbing, 1888,	540
cataphracta, Stebbing, 1888,	278, 352, 431
fragilis, Stebbing, 1888,	356, 394, 424
grimaldii, Stebbing, 1888,	1641

¹ Echinimana is a provisional name only.² The species of Proto printed in plain letters belong to the genus Phtisica, Slabber.³ Spence Bate, Brit. Mus. Catal. Amph. Crust., p. 169, gives this name to "Gammarus fimbriatus, Stimpson, MS.," but he says of it, "I do not feel quite satisfied that this species is distinct from P. pinguis."⁴ Limicolus, in the index to Stimpson's work.