

Rhachotropis— <i>continued.</i>			Siphonocetes— <i>continued.</i>	
<i>inflata</i> , Stebbing, 1888,	540		<i>pallidus</i> , G. O. Sars, 1882,	541
<i>kergueleni</i> , Stebbing, 1888,	955, pl. lxxxv		<i>typicus</i> , Krøyer, 1845,	{ 212, 217, 293, 302, 358 396, 430, 541, 1634}
<i>oculata</i> , Stebbing, 1888,	1644		<i>Whitei</i> , Sp. Bate, 1857,	275, 295, 430, 596
<i>Rhoea</i> —			<i>Sirenocyamus</i> ?—	
<i>Latreillii</i> , M.-Edw., 1828,	134		<i>Rhytinæ</i> , J. F. Brandt, 1847,	15, 218, 227, 405
<i>Sannazaria</i> —			<i>Socarnes</i> —	
<i>pallida</i> , O. and A. Costa, 1840,	183, 249		<i>bidenticulatus</i> , G. O. Sars, 1885,	{ 308, 466, 534, 568, 572 599, 600}
<i>Scamballa</i> (see <i>Orchestia</i>)—			<i>Kröyeri</i> , Boeck, 1872,	555
<i>Kuhliana</i> , Leach, MS., 1847,	222, 245		<i>ovalis</i> , Hock, 1882,	534, 572
<i>longicornis</i> , Leach, MS., 1847,	222, 246		{ <i>Vahli</i> , Boeck, 1870,	{ 161, 177, 214, 361, 393 466}
{ <i>megalophthalmos</i> , Brandt, 1851,	245		<i>Vahlia</i> , Lütken, 1875,	599, 1634
{ <i>megalophthalmus</i> , Leach, MS., 1847,	222			
<i>Sayana</i> , Leach, MS., 1847,	67, 222, 246			
<i>Tristensis</i> , Leach, MS., 1847,	222, 245			
<i>Schizoscelus</i> —			<i>Socarnoides</i> —	
<i>ornatus</i> , Claus, 1879,	492, 591, 1504		<i>kergueleni</i> , Stebbing, 1888,	691, pl. xxv
<i>rapax</i> , Bovallius, 1887,	143, 591, 1503		<i>Sophrosyne</i> —	
<i>Schnehagenia</i> —			<i>murrayi</i> , Stebbing, 1888,	652, pl. xv
<i>rapax</i> , Claus, 1871,	406, 492, 590		<i>Sperchius</i> —	
<i>Schraderia</i> —			<i>Lucidus</i> , Rafinesque, 1820,	111, 123
<i>gracilis</i> , Pfeffer, 1888,	1653		<i>Spinifer</i> —	
<i>Scia</i> (see <i>Clydonia</i> and <i>Tyro</i>)—			<i>flagelliformis</i> , Holbøll, MS., 1842,	198
<i>atlantica</i> , Stebbing, 1888,	558, 1270		<i>spinosissimus</i> , Holbøll, MS., 1842,	198
<i>borealis</i> , Stebbing, 1888,	592, 1270			
<i>clausii</i> , Stebbing, 1888,	558, 1272			
<i>cornigera</i> , Stebbing, 1888,	1273, pl. cxvi		<i>Squilla</i> —	
<i>ensicornis</i> , Prestandrea, 1833,	151, 183, 249, 1271		<i>acaudata</i> , Gronov, 1760,	19, 23
<i>longipes</i> , Stebbing, 1888,	265, 1270		<i>Balæni</i> , De Geer, 1778,	44
<i>marginata</i> , Stebbing, 1888,	558, 1272		<i>cauda nulla</i> ?, Gronov, 1762,	17, 23, 24
<i>pacifica</i> , Stebbing, 1888,	587, 1270		<i>fluviatilis</i> , Rösel, 1755,	{ 18, 21, 22, fig. 8, 359 1621}
<i>Tullbergii</i> , Stebbing, 1888,	558, 1270		<i>fluviatilis</i> , Merlet,	123
<i>Scopelocheirus</i> —			<i>lobata</i> , O. F. Müller, 1776,	{ 42, 46, 91, 180, 1618 1645}
{ <i>breviatus</i> , Sp. Bate, 1856,	290		<i>mantis (Amboinensis)</i> , Seba, 1758,	18
{ <i>crenatus</i> , Sp. Bate, 1857,	290, 292		<i>Pulex</i> , De Geer, 1778,	44, 113, 222
<i>Seba</i> —			<i>quadrilobata</i> , O. F. Müller, 1788,	53, 85
<i>innominata</i> (author unknown), {	18, 334, 560		<i>saltatrix</i> , Klein, 1743,	18, 185
Sp. Bate, 1862,			<i>ventricosa</i> , O. F. Müller, 1776,	42, 53, 86, 91, 1718
<i>Saundersii</i> , Stebbing, 1875,	451, 550, 783, pl. xlix			
<i>Simorhynchotus</i> —			<i>Stebbingia</i> —	
<i>antennarius</i> , Stebbing, 1888,	1572, pl. cc		<i>gregaria</i> , Pfeffer, 1888,	1653
<i>Lilljeborgi</i> , Stebbing, 1888,	1572		<i>Stegocephalus</i> —	
<i>Simorhynchus</i> (now <i>Simorhynchotus</i>)—			<i>ampulla</i> , ¹ Bell and Westwood, 1855,	{ 36, 214, 281, 355, 394 450, 531, 599, 729 1626, 1634}
<i>antennarius</i> , Claus, 1871,	406, 493, 590		<i>auratus</i> , G. O. Sars, 1882,	538
<i>Lilljeborgi</i> , Bovallius, 1887,	590		<i>Christianiensis</i> , Boeck, 1870,	{ 394, ² 538, ² 729, 1650}
<i>rapax</i> , Claus, 1887,	493		<i>gibbosus</i> , G. O. Sars, 1882,	538
<i>Siphonocetes</i> —			<i>inflatus</i> , ³ Krøyer, 1842,	{ 198, 214, 217, 271, 301 355, 599, 600, 729 pl. cxxxvii, A}
<i>Colletti</i> , Boeck, 1870,	396, 466, 541		<i>kessleri</i> , Stuxberg, 1880,	523, 543, 599, 729
<i>crassicornis</i> , Sp. Bate, 1857,	293, 542, 571		<i>latus</i> , Haswell, 1880,	511, 564
<i>cuspidatus</i> , Metzger, 1871,	408, 421, 445			
<i>cuspidatus</i> , S. I. Smith, 1874,	487			
<i>dubius</i> , Sp. Bate, 1856,	290			
<i>Kröyeranus</i> , Sp. Bate, 1857,	293, 295			

¹ The figure which Westwood gives of "*Stegocephalus* (Krøyer.) *ampulla* (Phipps)" has the postero-lateral angle of the third pleon-segment acute and a little upturned, but otherwise it closely agrees with Krøyer's *Stegocephalus inflatus*. Bell gives "*Stegocephalus Ampulla*, Krøyer," in the synonymy by mistake for *Stegocephalus inflatus*. In the definition of *Stegocephalus* (see p. 198) Krøyer says, "Pedes quinti parvis pedibus tertii quartique parvis structura et directione similes," a peculiarity at which he himself expresses his surprise, *Naturh. Tidskrift*, ser. 2, Bd. 1. p. 528, but Dana and Boeck are no doubt right in regarding the statement as an error.

² On p. 394, line 4, and p. 538, line 42, for *Christianensis* read *Christianiensis*.

³ The statement that this species is the same as *Cancer ampulla*, Phipps, has lately been corrected by Hansen, see pp. 599, 729.