

*Atlanta mediterranea*, Costa.

1852. *Atlanta mediterranea*, Costa, Fauna Reg. Napoli, Eteropodi, p. 4, pl. i. figs. 1a, A.B.C.

*Habitat.*—Bay of Naples.

This appears to be a large species, supposing that Costa's figure (1a) represents the natural size. The shell is peculiar on account of the keel extending to the aperture as in *Oxygyrus keraudrenii*, and not embracing the whorls of the spire as is the case in *Atlanta peronii*.

*Atlanta violacea*, Gould.

1852. *Atlanta violacea*, Gould, U.S. Explor. Exped., vol. xii. p. 493; Atlas, figs. 599–599b.

1862. „ „ *id.*, Otia, Conch., p. 236.

*Habitat.*—Mid Atlantic (Gould).

*Atlanta tessellata*, Gould.

1852. *Atlanta tessellata*, Gould, U.S. Explor. Exped., vol. xii. p. 494; Atlas, figs. 600–600b.

1862. „ „ *id.*, Otia, Conch., p. 236.

*Habitat.*—Mid Atlantic (Gould).

*Atlanta primitia*, Gould.

1852. *Atlanta primitia*, Gould, U.S. Explor. Exped., vol. xii. p. 491; Atlas, fig. 597.

1862. „ „ *id.*, Otia, Conch., p. 235.

*Habitat.*—Equatorial Atlantic (Gould).

*Atlanta cunicula*, Gould.

1852. *Atlanta cunicula*, Gould, U.S. Explor. Exped., vol. xii. p. 492; Atlas, figs. 598–598c.

1852. „ „ *id.*, Otia, Conch., p. 236.

*Habitat.*—Pacific Ocean, west of the Sandwich Islands (Gould).

*Atlanta souleyeti*, Smith.

1852. *Atlanta lamanonii*, Eydoux and Souleyet, Voy. "Bonite," Zool., vol. ii. p. 371; Atlas (1841), pl. xviii. figs. 30–37.

1850. „ „ Gray, Fig. Moll. Anim., pl. cexl. fig. 3 (copy of Eydoux and Souleyet).

*Habitat.*—Atlantic Ocean.