STATION 78.

Aphroditaster gracilis, n.g., n.sp. Obtained at no other locality. Only species of the genus.

Pentagonaster lepidus, n.sp. Obtained at no other locality.

OPHIUROIDEA (Lyman, Zool. pt. 14).

Ophiernus vallincola, n.g., n.sp. Obtained also at Stations 146 and 156, 1375 and 1975 fathoms. Only species of the genus.

Ophiocten hastatum, n.sp. Obtained also at Stations 146 and 168, 1375 and 1100 fathoms.

Amphiura otteri, Ljungman. Obtained also at Stations 45, 50, and 76.

ECHINOIDEA (Agassiz, Zool. pt. 9).

Salenia varispina, Agassiz. For distribution see Station 23.

Phormosoma uranus, Wyville Thomson, n.sp. Obtained also at Station VI.

HOLOTHURIOIDEA (Théel, Zool. pt. 39).

Holothuria lactea, n.sp. Two fragmentary specimens; obtained also at Station 169, 700 fathoms.

OSTRACODA (Brady, Zool. pt. 3).

Cythere dictyon, n.sp. Widely distributed (see Station 24).

irpex, n.sp. Obtained also at Stations 73 and 335.

Cypridina gracilis, n.sp. Obtained at no other locality.

CIRRIPEDIA (Hoek, Zool. pt. 25).

Scalpellum acutum, n.sp. One specimen; obtained also at Station 170, 520 to 630 fathoms.

AMPHIPODA (Stebbing, Zool. pt. 67).

Amaryllis haswelli, n.sp. One specimen; obtained at no other locality.

Isopoda (Beddard, Zool. pt. 48).

Ischnosoma spinosum, n.sp. One specimen; obtained at no other locality.

Leiopus leptodactylus, n.g., n.sp. Numerous specimens; obtained at no other locality. Only species of the genus.

Anomura (Henderson, Zool. pt. 69).

Lithodes agassizii, Smith. Two specimens; obtained at no other locality by the Challenger. Recorded from east coast of United States.