KERGUELEN.

- Nymphon brevicaudatum, Miers. Numerous specimens (25 to 120 fathoms); obtained at no other locality by the Challenger. Recorded from Kerguelen.
- Colossendeis robusta, n.sp. One specimen (120 fathoms); obtained at no other locality.
 - megalonyx, n.sp. Two or three specimens (120 fathoms); obtained also at Stations 313 and 314, 55 and 70 fathoms.

LAMELLIBRANCHIATA (Smith, Zool. pt. 35).

Neæra kerguelenensis, n.sp. (120 fathoms); obtained at no other locality.

Thracia meridionalis, n.sp. (20 to 60 fathoms); obtained also at Station 145.

Anatina elliptica, King and Broderip, juv. (15 to 28 fathoms); obtained at no other locality by the Challenger. Recorded from Kerguelen and New South Shetland.

Davila (?) umbonata, n.sp. (25 fathoms); obtained also at Station 145.

Kellia suborbicularis (Montagu). Two specimens (28 fathoms); obtained at no other locality by the Challenger. Recorded from North Atlantic.

" nuculina, Martens. (20 to 120 fathoms); obtained also at Station 145.

,, cardiformis, n.sp. (28 fathoms); obtained at no other locality.

Cardita astartoides, Martens. (20 to 60 fathoms); obtained also at Station 150, 150 fathoms.

Yoldia isonata, Martens. (15 to 110 fathoms); obtained at no other locality by the Challenger.

" subæquilateralis, Smith. (28 fathoms); obtained at no other locality by the Challenger.

Malletia gigantea, Smith. (Shallow water to 60 fathoms); obtained at no other locality by the Challenger.

Mytilus magellanicus, Chemnitz. (Shore); obtained also at Station 315, 12 fathoms, and Fiji. Recorded from New Zealand.

kerguelensis, n.sp. (Shore); obtained at no other locality.

Modiolarca trapezina (Lamarck). (28 fathoms); obtained also at Stations 145, 311, and 315.

kerguelensis, n.sp. (25 fathoms); obtained at no other locality.

Lima (Limatula) pygmæa, Philippi. (28 to 60 fathoms); obtained also at Station 145.

Pecten clathratus, Martens. (120 fathoms); obtained at no other locality by the Challenger.