

GASTEROPODA :

- Aplysia depilans*, Linné.
Cerithium vulgatum, Bruguière.
 „ (*Lampania australis*) *australe*, Quoy
 and Gaimard.
Homalogryra atomus (Philippi).
Natica (Lunatia) grönlandica, Beck.
Odostomia rissoïdes, Hanley.
Puncturella noachina (Linné).
Ringicula doliaris, Gould.
Rissoa (Onoba) arenaria (Mighels).
Trochus (Photinula) tæniatus (Sowerby).

CEPHALOPODA :

- Argonauta argo*, Linné.
Eledone verrucosa, Verrill.
Rossia sublevis, Verrill.

POLYZOA :

- Alecto granulata*, M.-Edwards.
Chorizopora bronniartii (Audouin).
Cribrilina radiata (Moll).
Crisia eburnea (Linné).
Diachoris magellanica, Busk.
Diastopora patina (Lamarck).
Escharoides verruculata (Smitt).
Frondipora verrucosa (Lamouroux).

The following species, taken by the Challenger to the *north* of the tropics, is recorded from the south :—

POLYZOA :

- Idmonea irregularis*, Meneghini.

The following species, not taken by the Challenger, are stated in the Reports to have been procured only in the north and south :—

Hornera lichenoides (Linné).

„ *violacea*, Sars.

Idmonea marionensis, Busk.*Lichenopora hispida* (Fleming).*Membranipora crassimarginata*
 (Hincks).*Pustulopora deflexa* (Smitt).

„ *proboscidea*, M.-Edwards.

„ *proboscidioides* (Smitt).

Salicornaria tenuirostris, Busk.*Serupocellaria pilosa* (Audouin).*Tubulipora fimbria*, Lamarck.

„ *flabellaris* (Fabricius).

BRACHIOPODA :

Platydia anomioides (Scacchi).

TUNICATA :

Boltenia legumen, Lesson.*Leptoclinum tenue*, n.sp.*Microcosmus polymorphus*, Heller.*Styela gyrosa* (Heller).

FISHES :

Acanthias blainvilli, Risso.*Halargyreus johnsoni*, Günther.*Notacanthus bonapartii*, Risso.*Scopelus macrolepidotus* (Johnston).*Stomias boa*, Risso.