

Malachra radiata, Linn.

Malachra radiata, Linn., Syst. Veg., p. 518; DC., Prodr., i. p. 440; Griseb., Fl. Brit. W. Ind., p. 81.

FERNANDO-NORONHA. *Moseley*.

Widely dispersed in Tropical America.

STERCULIACEÆ.

Waltheria americana, Linn.

Waltheria americana, Linn., Sp. Pl., ed. 1, p. 637 (including W. Indies, Linn.); Griseb., Fl. Brit. W. Ind., p. 95.

FERNANDO-NORONHA. *Darwin*; *Moseley*.

Generally spread in warm countries.

GERANIACEÆ.

Oxalis noronhae, Oliver, n. sp.

Frutex erectus, 1–2 pedalis, vel tortuosus, nodosus et humilior, pilis simplicibus vel ramosis vestita. *Folia alterna*, longe petiolata, 3-foliolata; foliola ovato-elliptica, 1–2 poll. longa, acuminata, obtusiuscula, subtus pubescentia, supra glabrescentia, lateralia obliqua, brevissime petiolata; petiolus gracilis, semiteres, supra canaliculatus, undique pilosulus 1–2½ poll. longus. *Flores pauci*, conferti, subumbellati, breviter pedicellati, vix 6 lineas longi, pedunculis quam folia saepius longioribus; calyx pilosus vel setulosus, sepalis oblongo-lanceolatis obtusiusculis; petala lutea, obtusa, sepalis subdupo longiora; stamina basi brevissime coalita; ovarium glabrum, profunde 5-lobum, lobis dorso tricostatis, apice deinde et breviter obtuse productis; ovula solitaria.—Hook., Ic. Pl., xiiii. p. 21, t. 1226. Descriptio parum mutata.

FERNANDO-NORONHA, RAT ISLAND, and ST MICHAEL'S MOUNT.—Without special locality
—*C. Darwin*; *Moseley*.

"Allied to *Oxalis insipida*, St. Hil., and in facies near Gardner's *Oxalis* (No. 2080), collected in Piauhy."—D. Oliver, *loc. cit.*

AMPELIDEÆ.

Vitis sicyoides, Baker.

Vitis sicyoides, Baker in Mart. Fl. Bras., xiv. pars 2, p. 202.

Cissus sicyoides, Linn., Sp. Pl., ed. 2, p. 170; Griseb., Fl. Brit. W. Ind., p. 102.

FERNANDO-NORONHA. *Moseley*.

General in Tropical and Subtropical America.