

Fragments only, and doubtfully belonging to this species.

West Indies; South America; Africa; Mascarene Islands; Southern Asia; Australia; New Zealand; and the Antarctic Islands.

ROCCELLEI.

Roccella tinctoria, DC.

Roccella tinctoria, DC.; Nyl., Synop. Lich., i. p. 258; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 155, et in Melliss's St Hel., p. 373.

ST HELENA. *Melliss.*

Canary Islands; Senegambia; Cape of Good Hope; East India; Central and South America; and also in the islands of the Mediterranean Sea; but here, according to Nylander, represented by a form approaching *Roccella phycopsis*.

Roccella phycopsis, Acharius.

Roccella phycopsis, Acharius; Nyl., Synop. Lich., i. p. 259; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 155, et in Melliss's St Hel., p. 373.

ST HELENA. *Burchell; Melliss.*

Mediterranean; Western Europe; Canaries; Ascension; Peru; and Madagascar.

Roccella fuciformis, Acharius.

Roccella fuciformis, Acharius; Nyl., Synop. Lich., i. p. 260; Leighton in Melliss's St Hel., p. 373.

ST HELENA. *Melliss.*

Mediterranean; Western Europe; Canaries; South America; Western and South Africa; and Mauritius.

USNEEI.

Usnea barbata, Fries.

Usnea barbata, Fries; Nyl., Synop. Lich., i. p. 267; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 374 (formæ: *florida*, *hirta*, *ceratina*, et *articulata*).

ST HELENA. *Burchell; Melliss.*

Generally diffused.

[*Usnea leucoxantha*, Bory in Duperr. Voy. "Coquille," Bot. Crypt., p. 241, St Helena, Durville, is unknown to us.]

Usnea laevis, Nyl.

Usnea laevis, Nyl., Synop. Lich., i. p. 271; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 271, et in Melliss's St Hel., p. 374.

ST HELENA. *Burchell.*