

# ENUMERATION OF THE PLANTS.

## PHANEROGAMÆ.

### CRUCIFERÆ.

#### *Pringlea antiscorbutica*, R. Br.

*Pringlea antiscorbutica*, R. Br. in Hook. f., Fl. Antarct., p. 239 t. 90; Oliver in Journ. Linn. Soc. Lond., xiv. p. 389.

HEARD ISLAND. *Moseley*.

Kerguelen, Crozets, and Marion Islands.

The Heard Island form is a very dwarf, though stout one, owing, doubtless, to a less fertile soil.

### CARYOPHYLLÆ.

#### *Colobanthus kerguelensis*, Hook. f.

*Colobanthus kerguelensis*, Hook. f., Fl. Antarct., p. 249 t. 92, et in Phil. Trans. Roy. Soc. Lond., clxviii. p. 19; Oliver in Journ. Linn. Soc. Lond., xiv. p. 389.

HEARD ISLAND. *Moseley*.

Elsewhere only known from Kerguelen Island.

### HALORAGEÆ.

#### *Callitriche verna*, Linn., subsp. *obtusangula*, Hook. f.

*Callitriche verna*, Linn., subsp. *obtusangula*, Hook. f. in Phil. Trans. Roy. Soc. Lond., clxviii. p. 20.

*Callitriche obtusangula*, Le Gall.; Hegelm., Monogr. Callitr., p. 54.

*Callitriche verna*, Linn., Sp. Pl., ed. 2, p. 6; Hook. f., Fl. Antarct., p. 272.

*Callitriche antarctica*, Engelm.; Oliver in Journ. Linn. Soc. Lond., xiv. p. 389.

HEARD ISLAND. *Moseley*.

Very widely dispersed in the temperate regions of both hemispheres.

### UMBELLIFERÆ.

#### *Azorella selago*, Hook. f.

*Azorella selago*, Hook. f., Fl. Antarct., p. 284, t. 99; Oliver in Journ. Linn. Soc. Lond., xiv. p. 389.

HEARD ISLAND. *Moseley*.

In many of the islets from Cape Horn eastward to Macquarie Island.