

Corallina berteri, Mont.

Corallina berteri, Mont. in Harv. Nereis Australis, p. 103, 1847, et in Gay, Fl. Chil., viii. p. 318 (1852, *berteroana*); Dickie in Journ. Linn. Soc. Lond., xv. p. 452; J. G. Agardh, Sp. Alg., i. p. 566.

JUAN FERNANDEZ. *Moseley*.

Melobesia pustulata, Lamour.

Melobesia pustulata, Lamour, Hist. Polyp. Flex., p. 315, t. 12; Kütz., Sp. Alg., p. 696; Harv., Nereis Australis, p. 110; Dickie in Journ. Linn. Soc. Lond., xv. p. 452.

JUAN FERNANDEZ. *Moseley*.

Atlantic, Mediterranean, and Adriatic, and in the Pacific on the coast of Norfolk Island.

Gigartina contorta, Bory.

Gigartina contorta, Bory in Duperr. Voy. "Coquille," Bot. Crypt., p. 153; J. G. Agardh, Sp. Alg., iii. p. 190; Gay, Fl. Chil., viii. p. 335; Dickie in Journ. Linn. Soc. Lond., xv. p. 452.

JUAN FERNANDEZ. *Moseley*.

The only previous record of this species is Concepcion (D'Urville).

Ceramium rubrum, Huds.

Ceramium rubrum, Huds., Fl. Angl., p. 600; J. G. Agardh, Sp. Alg., iii. p. 100; Hook. f., Handb. Fl. N. Zeal., p. 707; Gay, Fl. Chil., viii. p. 356; Dickie in Journ. Linn. Soc. Lond., xv. p. 452.

JUAN FERNANDEZ. *Moseley*.

A common and variable seaweed inhabiting the temperate seas of both hemispheres.

Centroceras micracanthum, Kütz.

Centroceras micracanthum, Kütz. in Linnæa, 1842, p. 741, et Sp. Alg., p. 688; Dickie in Journ. Linn. Soc. Lond., xv. p. 452 (*sphalmate micranthum*).

JUAN FERNANDEZ. *Moseley*.

Ptilota parva, Dickie.

Pinnis oppositis difformibus, una minore simplici, subulata, altera majore, pinnulata, pinnulis subulatis obtusis, vel subconicis, omnibus cellulosis. Sphærosporis—? Favellis—? (Dickie in Journ. Linn. Soc. Lond., xv. p. 452).

JUAN FERNANDEZ.—A few specimens mixed with *Centroceras*—*Moseley*.