

Polyporus (Inodermei) floridanus, Berkl.

Polyporus (Inodermei) floridanus, Berkl. in Ann. Nat. Hist., x. p. 376; et in Journ. Linn. Soc. Lond., xvi. p. 43.

ARROU.—British North America, southward through the West Indies and Mexico, to Bolivia and Brazil.

Trametes occidentalis, Fries.

Trametes occidentalis, Fries; Klotzsch in Linnæa, viii. p. 486; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

ARROU; KI.—Generally diffused in tropical and subtropical regions.

Trametes cognatus, Berkl.

Trametes cognatus, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

Pileo dimidiato sulcato-zonato subtiliter scabroso-tomentoso postice glabrescente ochraceo; hymenio plano, poris punctiformibus acie subtiliter tomentosis.

Pileus $1\frac{1}{2}$ inch across, 1 inch long; pores $\frac{1}{160}$ inch in diameter. Externally resembling *Trametes mülleri*, Berkl.; but the pores are very different.

ARROU.

Trametes sanguineus, Klotzsch.

Polyporus persoonii, Fries; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 45.

KI.—India; Ceylon; Burmah; Samoa and Friendly Islands; Gold Coast; Brazil.

Trametes bicolor, Berkl.

Trametes bicolor, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

Pileo reniformi lobato postice sanguineo, antice pallido ruguloso; hymenio umbrino; poris e subhexagonis elongatis; contextu pallido.

Pileus 2 inches across, 1 inch long; pores $\frac{1}{60}$ inch in diameter.

ARROU.

Hexagona arata, Berkl.

Hexagona arata, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

Tenuis rigidiuscula imbricata dimidiata v. flabelliformis regulariter sulcato-zonata scabroso-tomentosa umbrinella; hymenio e luteo-umbrino cinnamomeo; poris dentatis.

Pores $\frac{1}{48}$ inch in diameter. Allied to *Hexagona sericea*.

ARROU.