

Premna taitensis, Schauer.

Premna taitensis, Schauer ; Seem., Fl. Vit., p. 186, t. 43 ; Nadeaud, Enum. Pl. Tahiti, p. 57.
Scrophularioides arborea, Forst., Fl. Ins. Austr. Prodr., p. 91.

ADMIRALTY ISLANDS.—This is common in the Fijis, in the Ellice group, Samoa, and eastward to the Marquesas.

INCOMPLETÆ.

AMARANTACEÆ.

Cyathula prostrata, Blume.

Cyathula prostrata, Blume ; Seem., Fl. Vit., p. 199 ; Baker, Fl. Maurit., p. 269.

ADMIRALTY ISLANDS.—An annual herb, widely spread in Tropical Asia, Africa, and in Polynesia.

MYRISTICÆÆ.

Myristica glabra, Blume, var.

Myristica glabra, Blume, var. ; Miq., Fl. Ind. Bat., i. 2, p. 65.

ADMIRALTY ISLANDS.—From Silhet southward to Sumatra and Java.

LAURINEÆ.

Hernandia peltata, Meissn.

Hernandia peltata, Meissn. ; Seem., Fl. Vit., p. 204 ; Nadeaud, Enum. Pl. Tahiti, p. 48.

ADMIRALTY ISLANDS.—Tropical Asia, Africa, and Australia, and eastward in Polynesia as far as Tahiti.

THYMELÆACEÆ.

Phaleria blumei, Benth.

Phaleria blumei, Benth., Fl. Austr., vi. p. 38.
Drynispermum blumei, Dcne. ; Miq., Fl. Ind. Bat., i. 1, p. 885.

ADMIRALTY ISLANDS.—Malayan Archipelago and North-eastern Australia. The genus is restricted to Tropical Asia, and the southern area of the present species.

EUPHORBIACEÆ.

Euphorbia atoto, Forst.

Euphorbia atoto, Forst. ; Seem., Fl. Vit., p. 216 ; Nadeaud, Enum. Pl. Tahiti, p. 72 ; Benth., Fl. Austr., vi. p. 46.

ADMIRALTY ISLANDS.—India, Ceylon, Malayan Archipelago, and North-eastern Australia, and in Polynesia it extends eastward to Tahiti, and northward to the Marianne Islands, and closely allied species occur in Pitcairn, Elizabeth, and the Sandwich Islands.