

Station 163, April 4, 1874; lat. $36^{\circ} 56'$ S., long. $150^{\circ} 30'$ E.; depth, 2200 fathoms; bottom temperature, $0^{\circ} 7$ C.; bottom, red clay. Trawled in 120 fathoms,¹ off Twofold Bay.

Polycarpa radicata.

Ascidia cylindracea.

Port Jackson, Australia; 17th April to 23rd May 1874; 2 to 20 fathoms.

Molgula forbesi.

Cynthia cerebriformis.

irregularis.

complanata.

Styela gyrosa.

exigua.

Polycarpa tinctor.

viridis.

longisiphonica.

radicata.

Ascidia pyriformis.

Canterbury, New Zealand.

Boltenia pachydermatina.

Between New Zealand and the Fiji Islands the following Tunicata were collected:—

Station 170, July 14, 1874; lat. $29^{\circ} 55'$ S., long. $178^{\circ} 14'$ W.; depth, 520 fathoms; bottom temperature, $6^{\circ} 0$ C.; bottom, rock.

Culeolus wyville-thomsoni.

Kandavu, the Fiji Islands; July 27, 1874.

Cynthia pallida.

Between the Fiji Islands and Hong Kong, China, the following Tunicata were dredged:—

Station 186, September 8, 1874; lat. $10^{\circ} 30'$ S., long. $142^{\circ} 18'$ E.; depth, 8 fathoms; bottom, coral sand.

Cynthia arenosa.

formosa.²

Station 188, September 10, 1874; lat. $9^{\circ} 59'$ S., long. $139^{\circ} 42'$ E.; depth, 28 fathoms; bottom, mud.

Microcosmus helleri.

¹ The two Ascidians are almost certainly from this depth.

² This species is labelled "Torres Straits, 3 to 11 fathoms."