

PLATE XIII.

Figs. 1 and 2. *Culeolus perlatus*, Suhm.

Figs. 3 and 4. *Culeolus moseleyi*, Herdman.

Figs. 5 and 6. *Culeolus wyville-thomsoni*, Herdman.

Figs. 7-10. *Fungulus cinereus*, n. sp.

Fig. 1. Part of the peduncle of *Culeolus perlatus*, showing the brown trabeculae : enlarged.

Fig. 2. Part of the surface of the test, showing the papillæ ; enlarged.

Fig. 3. Part of the endostyle of *Culeolus moseleyi*, showing the spicules (sp.), the clear edges (w.e.), and the central opaque area ; magnified (Hartn., obj. 4).

Fig. 4. The nerve ganglion of *Culeolus moseleyi*, and the patch of spicules in the mantle over it ; magnified (Hartn., obj. 4).

Fig. 5. A small part of the branchial sac of *Culeolus wyville-thomsoni*, seen from the outside ; magnified (Hartn., obj. 4).

Fig. 6. Part of the endostyle of *Culeolus wyville-thomsoni*, showing the clear edge (w.e.) the calcareous spicules (sp.), and the central area of opaque brown bands ; magnified (Hartn., obj. 4).

Fig. 7. *Fungulus cinereus*, from the right side ; natural size.

Fig. 8. The branchial aperture ; enlarged.

Fig. 9. Part of the branchial sac, seen from the inside, showing one of the branchial folds (br.f.) ; magnified (Hartn., obj. 4).

Fig. 10. Part of the endostyle, showing the clear edges (w.e.), and the central opaque area. The left side of the figure shows the sinuses injected with blood corpuscles ; magnified (Hartn., obj. 4).