

PLATE LXXVI.

SAGRINA.—RAMULINA.—NUBECULARIA.

	Diam.	Page
Figures 1-3.— <i>Sagrina dimorpha</i> , Parker and Jones, . . .	× 60	582
Figs. 1, 2, 3. Lateral aspect; <i>b</i> , oral aspect.		
„ 4-10.— <i>Sagrina virgula</i> , Brady,		583
Figs. 4-7. Typical shallow-water form,		
Figs. 8-10. Monomorphous thicker-shelled specimens		
from deeper water,		
	× 70	
	× 50-70	
„ 11-16.— <i>Nubecularia divaricata</i> , Brady,	× 70	136
Figs. 11-14. Lateral aspect.		
Figs. 15, 16. Oral end of the test, showing the grooved		
tubular neck and everted lip.		
„ 17-19.— <i>Sagrina</i> (?) <i>tessellata</i> , n. sp.,		585
Figs. 17, 18. Lateral aspect; <i>b</i> , oral aspect,		
Fig. 19. Terminal segment more highly magnified,		
	× 75	
	× 150	
„ 20, 21.— <i>Sagrina</i> (?) <i>annulata</i> , n. sp.,		586
Fig. 20. Lateral aspect; <i>b</i> , oral aspect,		
Fig. 21. Terminal segment more highly magnified,		
	× 60	
	× 100	
„ 22-28.— <i>Ramulina globulifera</i> , Brady,	× 35	587