

PLATE CVI.

PULVINULINA.—ROTALIA.

		Diam.	Page
Figure	1.— <i>Pulvinulina berthelotiana</i> , d'Orbigny, sp.,	× 35	701
	<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
,,	2, 3.— <i>Pulvinulina lateralis</i> , Terquem, sp.,	× 35–40	689
	<i>a,a</i> , Superior aspect; <i>b,b</i> , inferior aspect; <i>c,c</i> , peripheral aspect.		
,,	4.— <i>Pulvinulina oblonga</i> , Williamson, sp.,	× 40	688
	<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
,,	5.— <i>Pulvinulina auricula</i> , Fichtel and Moll, sp.,	× 40	688
	<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
,,	6, 7.— <i>Pulvinulina hauerii</i> , d'Orbigny, sp.,	× 40–50	690
	<i>a,a</i> , Superior aspect; <i>b,b</i> , inferior aspect; <i>c,c</i> , peripheral aspect.		
,,	8.— <i>Pulvinulina oblonga</i> , var. <i>scabra</i> , nov.,	× 50	689
	<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
,,	9.— <i>Rotalia papillosa</i> , n. sp.,	× 30	708
	<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		