

furrowed longitudinally. Upper latus large, hexagonal. Infra-median latus large, wine-glass shaped. Scutum almost triangular. Peduncle cylindrical, with minute and very prominent scales.

This species is extremely characteristic through the form of its scutum and of its upper latus. It is represented by a single specimen only.

*Capitulum* of an elongate-oval shape, flat, consisting of thirteen valves only. The valves show distinct lines of growth, and are separated from each other by rather broad chitinous interspaces.

*Scutum* triangular, caused by the lateral margin being almost a prolongation of the short basal margin. Occludent margin arched. Tergal margin rather long, a triangular portion is described at the lateral extremity of the tergal margin by this margin, and by the uppermost part of the lateral margin. This triangular portion penetrates between the upper latus and the tergum, where it is received into a small excavation. Umbo of the scutum at the apex, which is slightly turned upwards.

*Tergum* triangular, with a not very long occludent margin. Carinal margin slightly arched, scutal margin slightly hollowed out. A triangular portion at the basal angle penetrates between the upper latus and the carina.

*Carina* not very strongly and simply bowed. Roof flat, not increasing much in width from the upper to the lower extremity; bordered by distinct ridges, hence the surface appears longitudinally furrowed. Sides at right angles with the roof, not very wide, but of the same width over the whole length of the carina.

*Upper latus* irregularly hexagonal, rather large. Its umbo is seated a little above the middle of the scutal margin.

*Rostral latus* irregularly quadrangular. Scutal margin almost straight, lateral margin describing an S-shaped line, rostral margin straight, basal margin straight also.

*Infra-median latus* having the form of a wine-glass. The foot is small, triangular, and the umbo is seated at the base of a much larger and wider portion.

*Carinal latus* irregularly quadrangular. Umbo at the foot of the carinal margin, slightly protuberant beyond the edge of the carina. Carinal margin hollowed out, upper margin short and oblique. Lateral margin hollowed out, basal margin well-developed.

Length of the capitulum, 15.5 mm.

*Peduncle* short, 5.5 mm., covered by very small and numerous calcareous scales, placed in not very regular rows. On some of the longitudinal rows thirteen or fourteen scales may be distinguished. The scales have very prominent and rounded free edges. The chitin of the peduncle may everywhere be seen between the scales.

Structure of the animal not studied; complementary male not observed.

The species was taken at Station 218, March 1, 1875; lat. 2° 33' S., long. 144° 4' E.; depth, 1070 fathoms; bottom temperature, 2° 1 C.; bottom, Globigerina ooze.