

II. *Marionia*, Vayssière.

- Marionia*, A. Vayssière, Sur un nouveau genre de la famille des Tritoniadés, Comptes rendus, t. lxxv., 1877, pp. 299-301.
- „ A. Vayssière, Description de la *Marionia Berghii*, Journ. de Conchyl., sér. 3, t. xix., 1879, pp. 106-118, pl. vii.
- „ R. Bergh, Beitr. zu einer Monogr. d. Gatt. *Marionia*, Vayss., Mittheil. d. zool. Stat. zu Neapel, Bd. iv., 1883, pp. 297-320, Taf. xxi.

Forma corporis ut in Tritoniis propriis ; limbus frontalis digitatus, digitis compositis numerosis.

Ventriculus lamellis solidis armatus.

The genus *Marionia* is readily distinguished from *Tritonia* by the form of the processes of the frontal veil, and by the armed condition of the stomach.

A number of species have been referred to this genus, many of which may possibly prove to be identical ; they are as follows :—

1. *Marionia blainvillea* (Risso).  
*Marionia berghii*, Vayss.  
Mediterranean.
2. *Marionia decaphylla* (Cantraine).  
Mediterranean.
3. *Marionia quadrilatera* (Schultz).  
Mediterranean.
4. *Marionia affinis*, Bergh.  
Mediterranean.
5. *Marionia tethydea* (delle Chiaje).  
Mediterranean.
6. *Marionia costæ* (Verany).  
Mediterranean.
7. *Marionia meyeri* (Verany).  
Mediterranean.
8. *Marionia acuminata* (O. G. Costa).  
Mediterranean.