

squamous suture, in the females and child the reverse was the case. The interzygomatic diameter was considerably below the interparietal. The mean interorbital diameter was 23.5, the maximum 26, the minimum 21. The bridge of the nose was slightly concavo-convex, the nasal bones varied in length from 21 to 25, and in greatest breadth from

TABLE X.—OAHU, SANDWICH ISLANDS, CHAL.

Brachycephali.

| Collection, | K. | S. | V. | Aa. | I*. | Ac. | Ah. |
|---------------------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|
| Age, | Ad. | Ad. | Ad. | 21 | Ad. | Ad. | Æt. 8 |
| Sex, | M. | M. | M. | M. | F. | F. | M? |
| Cubic capacity, | 1398 | 1675 | 1535 | 1430 | 1345 | 1495 | 1508 |
| Glabello-occipital length, | 176 | 184 | 182 | 174 | 175 | 178 | 164 |
| Ophryo-occipital " | 174 | 180 | 179 | 174 | 174 | 176 | 165 |
| Basi-bregmatic height, | 140 | 144 | 142 | 141 | 136 | 140 | 132 |
| Vertical Index, | 79 | 78 | 78 | 81 | 78 | 78 | 80 |
| Minimum frontal diameter, | 96 | 98 | 97 | 92 | 94 | 88 | 92 |
| Stephanic, " | 121 | 126 | 115 | 112 | 112 | 115 | 117 |
| Asterionic, " | 102 | 116 | 110 | 105 | 105 | 102 | 114 |
| Greatest breadth, | 146 _p | 155 _p | 148 _p | 150 _p | 146 _s | 146 _s | 152 _s |
| Cephalic Index, | 83 | 84 | 81 | 86 | 83 | 82 | 92 |
| Horizontal circumference, | 506 | 541 | 523 | 504 | 512 | 506 | 499 |
| Frontal longitudinal arc, | 126 | 140 | 128 | 128 | 125 | 127 | 124 |
| Parietal " | 136 | 107 | 124 | 128 | 125 | 126 | 117 |
| Occipital " | 112 | ... | 118 | 105 | 120 | 118 | 109 |
| Total " | 374 | ... | 370 | 361 | 370 | 371 | 350 |
| Vertical transverse arc, | 313 | 338 | 325 | 333 | 310 | 318 | 328 |
| Length of foramen magnum, | 29 | ... | 36 | 35 | 32 | 32 | 34 |
| Basi-nasal length, | 100 | 108 | 110 | ... | 100 | 104 | 89 |
| Basi-alveolar length, | 97 | ... | 110 | ... | 102 | 104 | 88 |
| Gnathic Index, | 97 | ... | 100 | ... | 102 | 100 | 99 |
| Interzygomatic breadth, | 129 | 142 | 138 | ... | 126 | 129 | 119 |
| Intermalar " | 115 | 124 | 119 | ... | 110 | 110 | 108 |
| Ophryo-alveolar length, | 88 | ... | 89 | ... | 85 | 86 | 73 |
| Naso-alveolar " | 69 | ... | 66 | ... | 63 | 63 | 57 |
| Facial Index, | 68 | ... | 64 | ... | 67 | 67 | 61 |
| Nasal height, | 48 | ... | 51 | ... | 47 | 50 | 42 |
| Nasal width, | 22 | ... | 23 | ... | 27 | 24 | 20 |
| Nasal Index, | 46 | ... | 45 | ... | 57 | 48 | 47 |
| Orbital width, | 37 | 38 | 39 | ... | 36 | 36 | 34 |
| Orbital height, | 35 | 35 | 31 | ... | 33 | 37 | 33 |
| Orbital Index, | 95 | 92 | 79.5 | ... | 92 | 103 | 97 |
| Palato-maxillary length, | 53 | ... | 58 | ... | 55 | 54 | 45 |
| Palato-maxillary breadth, | ... | ... | 65 | ... | 66 | ... | 59 |
| Palato-maxillary Index, | ... | ... | 112 | ... | 124 | ... | 131 |
| Lower Jaw. { Symphysial height, | 26 | ... | ... | ... | 28 | ... | ... |
| Coronoid " | 64 | ... | ... | ... | 60 | ... | ... |
| Condylloid " | 62 | ... | ... | ... | 53 | ... | ... |
| Gonio-symphysial length, | 86 | ... | ... | ... | 88 | ... | ... |
| Intergonial width, | 89 | ... | ... | ... | 86 | ... | ... |
| Breadth of ascending ramus, | 33 | ... | ... | ... | 35 | ... | ... |

* With pelvis.