

PLATE IV_A.

- Fig. 1. Bristles from the ventral group of the first foot of *Lætmonice producta*, × 350.
- Fig. 2. One of the stouter superior bristles showing minute chitinous nodules. From the second foot of the same, × 350.
- Fig. 3. Serrated bristle from the upper division of the ventral branch of the second foot in the same form, × 90.
- Fig. 4. Spinose bristle from the lower division of the ventral branch of the second foot of the preceding, × 90.
- Fig. 5. Tip of one of the lustrous brown dorsal spines (of medium size) of the same species, × 90.
- Fig. 6. One of the same with the chitinous sheaths or guards from the ninth foot, × 90.
- Fig. 7. Tip of a spine showing an alternate arrangement of the recurved teeth from the same species, × 55.
- Fig. 8. Tip of one of the fully formed ventral bristles of the foregoing, × 55.
- Fig. 9. Tip of one of the dark brown dorsal spines of *Lætmonice producta*, var. *wyvillei*, × 90.
- Fig. 10. Tip of one of the ventral bristles of the same form, × 55.
- Fig. 11. Portion of the distal region of one of the simple inner dorsal bristles of the same, showing the nodes, which are better marked than in *Lætmonice producta* proper, × 350.
- Fig. 12. Portion of one of the fine dorsal bristles overlapping the scales of *Lætmonice producta*, var. *benthaliana*, × 350.
- Fig. 13. Portion of the shaft (a short distance below that figured in Pl. V_A. fig. 10) of one of the dorsal spines of *Lætmonice japonica*, × 90.