

PLATE XIIA.

Fig. 1. Tip of dorsal bristle of <i>Lagisca darwini</i> ,	· · · · ·	× 800.
Fig. 2. Extremity of one of the longer ventral bristles of the same,	· · · · ·	× 800.
Fig. 3. One of the shorter ventral bristles of the foregoing,	· · · · ·	× 700.
Fig. 4. Extremity of a dorsal bristle of <i>Lagisca kermadecensis</i> ,	· · · · ·	× 350.
Fig. 5. One of the upper ventral bristles of the same in profile,	· · · · ·	× 350.
Fig. 6. Tip of one of the latter. Viewed antero-posteriorly,	· · · · ·	× 350.
Fig. 7. Dorsal bristle of <i>Eunoa mindanavensis</i> ,	· · · · ·	× 350.
Fig. 8. Tip of a ventral bristle of the same species,	· · · · ·	× 250.
Fig. 9. Smooth dorsal bristle of <i>Polynoë (Macellicephala) mirabilis</i> ,	· · · · ·	× 80.
Fig. 10. One of the ventral bristles of the foregoing,	· · · · ·	× 80.
Fig. 11. Tip of one of the latter,	· · · · ·	× 400.
Fig. 12. Tip of one of the dorsal bristles of <i>Polynoë synophtalma</i> ,	· · · · ·	× 250.
Fig. 13. Tip of a ventral bristle of the same species,	· · · · ·	× 250.
Fig. 14. One of the shorter dorsal bristles of <i>Lagisca hexactinellidæ</i> ,	· · · · ·	× 400.
Fig. 15. One of the superior (long) ventral bristles of the same,	· · · · ·	× 400.
Fig. 16. One of the inferior (shorter) ventral bristles,	· · · · ·	× 400.
Fig. 17. Tip of a ventral bristle of <i>Polynoë (Admetella) longipedata</i> ,	· · · · ·	× 210.
Fig. 18. Tip of an average ventral bristle, from about the twentieth foot of <i>Polynoë ocellata</i> ,	· · · · ·	× 800.
Fig. 19. Tip of one of the powerful bristles, from the fiftieth foot of the same. The contrast with the former is bold,	· · · · ·	× 800.