

2. *Marginella (Glabella) musica*, Hinds.

- Marginella musica*, Hinds, Proc. Zool. Soc. Lond., 1844, p. 73.
 " " Sowerby, Thes. Conch., pt. 7, p. 379, sp. 23, pl. lxxv. figs. 36, 37.
 " " Reeve, Conch. Icon., vol. xv. pl. ix. fig. 34.
 " " Weinkauff, Conch. Cab. (ed. Küster), p. 52, sp. 65, pl. ix. figs. 13, 16.
 " (*Glabella musica*, Tryon, Manual, vol. v. p. 22, pl. vi. fig. 95.

Station VIII. February 12, 1873. Lat. 28° 3' 15" N., long. 17° 27' W. Gomera, Canaries. 620 fathoms. Volcanic mud.

Station 142. December 18, 1873. Lat. 35° 4' S., long. 18° 37' E. Cape of Good Hope. 150 fathoms. Green sand. Bottom temperature 47°.

Habitat.—West Africa. 35 fathoms. Sand. (Hinds.)

The specimen from the Cape has the more elongated form of *Marginella diadochus*, Adams and Reeve, but if the species be really different the locality suits Hinds' species better than that of Adams and Reeve. Neither from the descriptions, however, nor from an examination of the shells, can I consider the species of Adams and Reeve as distinct, and Hinds' name has priority.

3. *Marginella (Glabella) turbinata*, Sowerby.

- Marginella turbinata*, Sowerby, Thes. Conch., pt. 7, p. 385, sp. 46, pl. lxxv. figs. 70, 71.
 " " Reeve, Conch. Icon., vol. xv. pl. xxii. fig. 122.
 " " Angas, Port Jackson Moll., Proc. Zool. Soc. Lond., 1867, p. 196, No. 60.
 " " Weinkauff, Conch. Cab. (ed. Küster), 1879, p. 86, sp. 115, pl. xvi. figs. 9, 12.
 " (*Glabella turbinata*, Tryon, Manual, vol. v. p. 23, pl. vii. figs. 4, 5.

April 17, 1874. Port Jackson, Sydney. 2 to 10 fathoms.

May 1864. Port Jackson, Sydney. 6 to 7 fathoms.

Habitat.—Port Jackson, Australia (Angas).

4. *Marginella (Glabella) pachia*,¹ n. sp. (Pl. XVI. fig. 5).

Station 185B. August 31, 1874. Lat. 11° 38' 15" S., long. 143° 59' 38" E. Off Raine Island, Cape York, North-East Australia. 155 fathoms. Coral sand.

Shell.—Small, obovate, with scarcely any spire, strong, porcellanous white, lustrous, with an excessively narrow mouth, a truncate but not emarginate varixed toothless outer lip, and four small folds on the sharp-pointed pillar. *Sculpture*: None but faint lines of growth. *Colour* white, more or less translucent. *Spire* just perceptibly raised, but visible rather by colour than form, the suture being a little more opaque than the rest of the whorls. *Apex* minutely mamillate. *Whorls* 4, small, just perceptibly convex on the spire; the last is tumid, with a roundly conical and produced base. *Suture* barely

¹ παχύς, fat.