

This species is much more attenuated in front than *Dolabrifera marmorea*, Pease, from the Sandwich Islands, which otherwise it much resembles in form and texture, whilst it is in sculpture much more delicate. *Dolabrifera maillardi*, Deshayes, from the island of Bourbon (see Moll. de Bourbon, p. 53, vol. vii. figs. 20–22), is much more regular in shape, more like the seed of our common plane (*Acer pseudo-platanus*), with a regular-shaped wing and a head or nucleus continuous with the body instead of, as here, a fleam-like blade and distinct handle.

Family PLEUROBRANCHIDÆ, Gray, 1840.

Genus *Umbrella*, Lamarck, 1812.

Umbrella mediterranea, Lamarck.

- Umbrella mediterranea*, Lamarck, Anim. s. vert., vol. vi. p. 343, and (ed. Desh.) vol. vii. p. 574, sp. 2.
 " " Deshayes, Encycl. méthod., vers, vol. iii. p. 663, sp. 2.
 " " Philippi, Enumeratio Moll. Sicil., vol. i. p. 113, vol. ii. p. 88, pl. vii. fig. 11.
 " " Cantraine, Malacol. Med., p. 92.
 " " Adams, Genera, vol. ii. p. 91, pl. lxi. fig. 3.
 " " Weinkauff, Conch. d. Mittelm., vol. ii. p. 179.
 " " Monterosato, Enumerazione, p. 53.

July 28–31, 1873. St Vincent, Cape Verde Islands. Harbour. 7 to 25 fathoms.

Habitat.—Mediterranean (Lamarck); Madeira (Watson).

The embryonic apex of this shell is glossy, of a rich chestnut brown. It is placed a little to the left side of the shell, and is directed backwards. Including the whole shell, there are barely two whorls, and these are sinistral.

Family SIPHONARIIDÆ, Gray, 1840.

Genus *Siphonaria*, Sowerby, 1824.

Species.

1. *Siphonaria cancer*, Reeve.

2. *Siphonaria (Liriola) tristensis*, Sow.

1. *Siphonaria cancer*, Reeve:

- Siphonaria cancer*, Reeve, Conch. Icon., vol. ix. pl. ii. fig. 7.
 " " Hutton, Cat. Mar. Moll. New Zealand, p. 55, No. 261*.
 " " v. Martens, Crit. List., p. 40.
 " " Hutton, Manual, p. 36.

June 26, 1874. D'Urville Island, New Zealand. Beach.

Habitat.—New Zealand (British Museum).