

STATION 246. July 2, 1875; lat. $36^{\circ} 10'$ N., long. $178^{\circ} 0'$ E.; depth, 2050 fathoms; bottom, Globigerina ooze; bottom temperature, $35^{\circ}.1$. Trawled.

Aristeus armatus (1 ♂).

STATION 250. July 9, 1875; lat. $37^{\circ} 49'$ N., long. $166^{\circ} 47'$ W.; depth, 3050 fathoms; bottom, red clay; bottom temperature, 35° . Trawled.

Benthesicymus pleocanthus (2 ♂). | *Gennadas parvus* (1).

STATION 254. July 17, 1875; lat. $35^{\circ} 13'$ N., long. $154^{\circ} 43'$ W.; surface temperature, 72° . Surface.

Aristeus (young).

North Pacific; July 20, 1875.

Sergestes ancylops.

STATION 256. July 21, 1875; lat. $30^{\circ} 22'$ N., long. $154^{\circ} 56'$ W.; depth, 2950 fathoms; bottom, red clay; bottom temperature, $35^{\circ}.2$. Dredged.

Sergestes armatus (1).

STATION 257. July 23, 1875; lat. $27^{\circ} 33'$ N., long. $154^{\circ} 55'$ W.; surface temperature, $76^{\circ}.5$. Surface.

Sergestes oculatus (1).

Off the Sandwich Islands.

Sergestes parvidens.

Sergestes ventridentatus.

Lucifer (young).

Lucifer reynaudii.

North Pacific, August 21, 1875.

Sergestes oculatus.

Honolulu; fresh-water rivers.

Atya bisulcata (100 ♀ ♂).

| *Bithynis grandimanus*.

Honolulu (reefs).

Alpheus laevis (1 ♂).

Off Honolulu; depth, 18 fathoms.

Alpheus prolificus (1 ♀).