

PLATE I.

| | Diam. | Page |
|---|-------|------|
| Fig. 1. <i>Dromia ciliata</i> , n. sp., | × 1 | 3 |
| <i>a.</i> Cephalic region (with the hairs removed on the left side), | × 1 | |
| <i>b.</i> Fronto-orbital region (with hairs removed), | × 1 | |
| <i>c.</i> Abdomen of male, | × 1 | |
| Fig. 2. <i>Cryptodromia japonica</i> , n. sp., | × 1 | 6 |
| <i>a.</i> Cephalic region, | × 2 | |
| Fig. 3. <i>Cryptodromia nodulifera</i> , n. sp., | × 2 | 8 |
| <i>a.</i> Cephalic region, | × 2 | |
| <i>b.</i> Abdomen of male, | × 2 | |
| Fig. 4. <i>Cryptodromia incisa</i> , n. sp., | × 1 | 10 |
| <i>a.</i> Cephalic region (with the hairs removed on the left side), | × 2 | |
| <i>b.</i> Abdomen of female, | × 1 | |
| Fig. 5. <i>Dromidia antillensis</i> , Stimpson, | × 1 | 12 |
| <i>a.</i> Cephalic region, | × 2 | |
| <i>b.</i> Abdomen of female, | × 1 | |
| Fig. 6. <i>Dromidia spongiosa</i> , Stimpson, | × 1 | 12 |
| <i>a.</i> Cephalic region, | × 2 | |
| <i>b.</i> Abdomen of male, | × 1 | |
| Fig. 7. <i>Eudromia frontalis</i> , n. gen. et sp., | × 1 | 14 |
| <i>a.</i> Cephalic region, | × 2 | |
| <i>b.</i> Left chelipede of female, | × 1 | |
| Fig. 8. <i>Pseudodromia latens</i> , Stimpson (with the fifth right leg drawn back), | × 1 | 16 |
| <i>a.</i> Cephalic region, | × 2 | |
| <i>b.</i> Abdomen of male, | × 1 | |