

Between 1500 and 2000 fathoms.

Family.	Genus and Species.	Station.	Depth.	Bottom.	Region.
Bicellariadæ,	<i>Kinetoskias cyathus</i> , . . .	VI.	1090 to 1525	gl. oz.	N. Atlantic.
	<i>Bugula reticulata</i> , var. <i>unicornis</i> , . . .	106	1850	gl. oz.	"
	* " " . . .	147	1600	di. oz.	S. Indian.
	" " <i>leontodon</i> , . . .	3	1525	h. gr.	N. Atlantic.
	" " <i>bicornis</i> , . . .	157	1950	di. oz.	S. Indian.
	" " <i>margaritifera</i> , . . .	323	1900	bl. m.	S. Atlantic.
	<i>Bicellaria infundibulata</i> , . . .	147	1600	di. oz.	S. Indian.
"	" "	156	1975	di. oz.	"
Cellulariadæ,	<i>Canda simplex</i> , . . .	44	1700	bl. m.	N. Atlantic.
	<i>Menipea clausa</i> , . . .	70	1675	gl. oz.	"
	<i>Cellularia crateriformis</i> , . . .	323	1900	bl. m.	S. Atlantic.
	<i>Nellia simplex</i> , . . .	87	1675	rock.	N. Atlantic.
Farciminariadæ,	<i>Farciminaria delicatissima</i> , . . .	13	1900	gl. oz.	"
	" "	14	1950	"	"
	" <i>gracilis</i> , . . .	70	1675	"	"
	" <i>magna</i> , . . .	153	1675	bl. m.	S. Indian.
	" " var. <i>armata</i> , . . .	323	1900	bl. m.	S. Atlantic.
	" <i>cribraria</i> , . . .	323	1900	"	"
Bifaxariadæ,	<i>Bifaxaria reticulata</i> , . . .	13	1900	gl. oz.	N. Atlantic.
	" <i>minuta</i> , . . .	70	1675	"	"
Salicornariadæ,	* <i>Salicornaria magnifica</i> , . . .	13	1900	"	"
	" "	323	1900	bl. m.	S. Atlantic.
	" "	157	1950	di. oz.	S. Indian.
Membraniporidæ,	<i>Foveolaria orbicularis</i> , . . .	147	1600	di. oz.	"
Catenariadæ,	<i>Catenaria bicornis</i> , . . .	280	1940	gl. oz.	S. Pacific.
Onchoporidæ,	* <i>Onchopora sinclairii</i> , . . .	157	1950	di. oz.	S. Indian.
Escharidæ,	* <i>Tessaradoma boreale</i> , . . .	13	1900	gl. oz.	N. Atlantic.
Idmoneidæ,	<i>Idmonea marionensis</i> , . . .	147	1600	di. oz.	S. Indian.