

Figure	PAGE
273. Diagram representing a Nurse-Stock of the Mushroom Coral (<i>Fungia</i>) and enlarged view of the scar left on the end of the stock when a young coral has become detached,	789
274. Portion of the hard coral skeleton of <i>Millepora nodosa</i> ,	791
275. System of zooids of <i>Millepora nodosa</i> in the expanded condition,	791
276. <i>Cryptohelia pudica</i> , M.-Edw. and Haime,	792
277. <i>Deltocyathus italicus</i> , Pourtalès,	793
278. <i>Deltocyathus italicus</i> . Stellate variety of Pourtalès,	793
279. <i>Deltocyathus magnificus</i> , Moseley,	794
280. <i>Odontocyathus coronatus</i> , Moseley,	794
281. <i>Stephanotrochus diadema</i> , Moseley	794
282. <i>Stephanotrochus nobilis</i> , Moseley,	794
283. <i>Flabellum alabastrum</i> , Moseley,	795
284. <i>Flabellum angulare</i> , Moseley,	795
285. <i>Flabellum apertum</i> , Moseley,	796
286. <i>Lophohelia prolifera</i> , M.-Edw. and Haime,	796
287. <i>Bathyactis symmetrica</i> , Moseley,	797
288. <i>Stephanophyllia complicata</i> , Moseley,	798
289. <i>Leptopenus hypocælus</i> , Moseley,	798
290. <i>Leptopenus hypocælus</i> , Moseley,	799
291. Tooth of <i>Carcharodon megalodon</i> , from 2385 fathoms,	805
292. Petrous and tympanic bone of <i>Ziphius cavirostris</i> , from 2335 fathoms,	806
293. Section of a Manganese Nodule, showing a tympanic bone of <i>Mesoplodon</i> in the centre, from 2600 fathoms,	807
294. <i>Oxyrhina</i> (<i>Oxyrhina trigonodon</i> , Agass.) tooth, from 2350 fathoms,	808
295. Black Spherule with Metallic Nucleus from 2375 fathoms, South Pacific,	811
296. Black Spherule with Metallic Nucleus from 3150 fathoms, Atlantic,	811
297. Spherule of Bronzite from 3500 fathoms, Central South Pacific,	812
298. A Lamella of the Spherule represented in fig. 297,	812
299. Cumberland Bay, Juan Fernandez,	822
300. } <i>Carinina</i> , n. gen.,	831
301. }	
302. <i>Pelagonemertes rollestoni</i> , Moseley,	832
303. <i>Keramosphæra murrayi</i> , Brady,	834
304. <i>Cymbalopora</i> (<i>Tretomphalus</i>) <i>bulloides</i> (d'Orbigny),	835
305. <i>Hastigerina pelagica</i> (d'Orbigny) (<i>murrayi</i> , Wyv. Thoms.), with floating apparatus,	836
306. <i>Hastigerina pelagica</i> (d'Orbigny) (<i>murrayi</i> , Wyv. Thoms.), from the surface,	837
307. <i>Pulvinulina menardii</i> (d'Orbigny), from the tropical deposits,	838
308. <i>Globigerina bulloides</i> , d'Orbigny, from the deposits, tropical regions of the Atlantic,	839
309. <i>Globigerina bulloides</i> , var. <i>triloba</i> (Reuss), from the surface,	840
310. <i>Orbulina universa</i> (d'Orbigny), from the deposits,	841
311. <i>Orbulina universa</i> (d'Orbigny), from the surface,	842
312. <i>Corycæus pellucidus</i> , Dana; male and female,	844
313. <i>Pontostratiotes abyssicola</i> , G. S. Brady,	845
314. <i>Lernæa abyssicola</i> , G. S. Brady,	846
315. <i>Cythere dictyon</i> , G. S. Brady,	847
316. <i>Krithe producta</i> , G. S. Brady	847
317. <i>Megalasma striatum</i> , Hoek,	852
318. <i>Scapellum eximium</i> , Hoek,	853
319. <i>Verruca sulcata</i> , Hoek,	854