

See Charts 31 and 35, and Diagram 16.

Number of Station.	Date.	Position.	Depth in Fathoms.	Temperature of the Sea-water (Fahr.).		Designation and Physical Characters.	CARBONATE OF CALCIUM.			
				Bottom	Surface		Per cent.	Foraminifera.	Other Organisms.	
Yokohama—continued.	280	1875 April 5	26 29 0 N. 137 57 0 E.	2425	35.5	68.5	RED CLAY, similar to the last, fine grained, coherent, chocolate coloured and plastic when wet.
	281	" 9	31 8 0 N. 137 8 0 E.	2250	35.2	64.0	BLUE MUD, blue-grey, fine grained, coherent, presenting no macroscopic elements, breaking up readily in water.	trace	Globigerinidæ.	...
	282	May 12	35 11 0 N. 139 28 0 E.	345	41.1	64.2	GREEN MUD, green-grey, earthy, slightly coherent, breaking up readily in water, containing shells. Residue dark green with brown tinge.	3.29	(1.00%), Globigerinidæ, <i>Pulvinulina</i> , (1.00%), Textularidæ, Rotalidæ.	(1.29%), Ostracodes, Echinoderm fragments, Coccoliths.
	283	" 17	34 39 0 N. 135 14 0 E.	8	...	62.3	BLUE MUD, light blue-grey, slightly coherent, breaking up with difficulty in water. Residue green.	11.32	A few <i>Rotalia</i> .	(11.32%), Gasteropods, Lamellibranchs, Ostracodes, Echinoderm fragments.
	283A	" 19	34 38 0 N. 135 1 0 E.	50	...	62.6	SAND.
Off Japan.	283B	" 26	34 18 0 N. 133 36 0 E.	15	...	66.3	BLUE MUDS, light blue-grey, coherent, breaking up with difficulty in water, plastic when wet. Residue dark blue-grey.	4.29	(1.00%), Miliolidæ, Rotalidæ.	(3.29%), <i>Dentalium</i> , Gasteropods, Lamellibranchs, Ostracodes, Echinoderm fragments.
	283C	" 28	34 18 0 N. 133 21 0 E.	12	59.9	66.8				
	284	June 3	32 31 0 N. 135 39 0 E.	2675	35.8	69.5	BLUE MUD, light blue-grey, slightly coherent, fine grained, breaking up readily in water, dark blue-grey and somewhat plastic when wet. Residue dark blue-grey.	trace	<i>Globigerina bulloides</i> , <i>Bulimina ovata</i>
285	" 4	34 7 0 N. 138 0 0 E.	565	38.1	73.0	GREEN MUD, light green-grey, slightly coherent, breaking up in water. Residue green.	[5.00]	(2.00%), Globigerinidæ, <i>Pulvinulina menardii</i> , (1.00%), <i>Bolivina textularioides</i> , Lagenidæ, <i>Truncatulina pygmaea</i> .	(2.00%) Ostracodes, Echinoderm fragments, Coccoliths, Rhabdoliths.	