See Chart 16, and Diagram 6.

Rio de la Plata to Tristan da Cunha—*continued.*

Number of Station.	Date.	Position.	Depth in Fathoms.	Temperature of the Sea-water (Fahr. Bottom	Temperature of the Sea-water (Fahr. Surface	Designation and Physical Characters.	Carbonate of Calcium. Per cent.	Foraminifera.	Other Organisms.
	1876	° ′ ″		°	°				
326	Mar. 3	37 3 0 S. 44 17 0 W.	2775	32·7	67·8	BLUE MUD, grey-brown, arenaceous, plastic, finely granular, drying into slightly coherent earthy masses. **Residue** grey-brown.	3·11	...	(3·11 %), small teeth of fish.
327	,, 4	36 48 0 S. 42 45 0 W.	2900	32·8	70·2	RED CLAY, grey-brown, finely granular, slightly coherent, earthy.	...	...	...
328	,, 6	37 38 0 S. 39 36 0 W.	2900	32·9	68·0	RED CLAY, grey-brown, plastic, unctuous, coherent when dry, finely granular, sublustrous streak.	...	...	...
329	,, 7	37 31 0 S. 36 7 0 W.	2675	32·3	64·5	RED CLAY, grey-brown, plastic, unctuous, homogeneous, drying into fine-grained coherent masses, sublustrous streak.	0·70	Fragments of pelagic Foraminifera.	...
330	,, 8	37 45 0 S. 33 0 0 W.	2440	32·7	64·2	RED CLAY, grey-brown, homogeneous, coherent, lustrous streak. **Residue** brown.	10·36	(6·00 %), fragments of Globigerinidæ, *Pulvinulina*.	(4·36 %), Ostracodes, one or two fragments of Echini spines, Coccoliths.
331	,, 9	37 47 0 S. 30 20 0 W.	1715	35·4	64·5	GLOBIGERINA OOZE, white with a slight rose tint, granular, pulverulent. **Residue** brown.	78·38 *	(60·00 %), Globigerinidæ, *Pulvinulina*. (3·00 %), Miliolidæ, Textularidæ, Lagenidæ, Rotalidæ.	(15·38 %), Ostracodes, Echini spines, Polyzoa, Coccoliths, Coccospheres, Rhabdoliths.
*332	,, 10	37 29 0 S. 27 31 0 W.	2200	34·0	64·0	GLOBIGERINA OOZE, grey with a rose tinge, homogeneous, slightly coherent. **Residue** brown.	65·67	(55·00 %), Globigerinidæ, *Pulvinulina*. (3·00 %), Miliolidæ, Textularidæ, Lagenidæ, Rotalidæ.	(7·67 %), fragments of Pteropods, Echini spines, Coccoliths, Coccospheres, Rhabdoliths.
333	,, 13	35 36 0 S. 21 12 0 W.	2025	35·3	67·0	GLOBIGERINA OOZE, white, slightly coherent. **Residue** red.	88·97	(75·00 %), Globigerinidæ, *Pulvinulina*. (3·00 %), Miliolidæ, Textularidæ, Lagenidæ, Rotalidæ.	(10·97 %), fragments of Pteropods, Ostracodes, Echini spines, Polyzoa, Coccoliths, Coccospheres, Rhabdoliths.

* See anal. 52.