

STATION 24.

Lophohelia prolifera, M.-Edwards and Haime. Small fragment; for distribution see Station 23.

Bathyactis symmetrica (Pourtales). Very widely distributed, both geographically and bathymetrically; obtained at Stations 24, 36, 56, 73, 78, 133, 147, 157, 181, 194, 195, 196, 218, 224, 241, 244, 299, 325, and 332, 32 to 2900 fathoms.

HYDROIDA (Allman, Zool. pt. 70).

Cryptolaria flabellum, n.sp. Obtained at no other locality.

Sertularia catena, n.sp. One specimen; obtained at no other locality.

CRINOIDEA (Carpenter, Zool. pt. 32).

Rhizocrinus lofotensis, Sars. Two specimens from 625 fathoms; obtained also at Stations 122 and 323, 400 and 1900 fathoms. Recorded from both sides of the North Atlantic ("Porcupine," "Lightning," "Knight Errant," and various American expeditions).

ASTEROIDEA (Sladen, Zool. pt. 51).

Brisinga cricophora, n.sp. Obtained at no other locality.

OPHIUROIDEA (Lyman, Zool. pt. 14).

Ophiozona nivea, Lyman. Obtained at no other locality by the Challenger.

Ophioglypha falcifera, Lyman. Obtained at no other locality by the Challenger.

„ *variabilis*, n.sp. (young?). Obtained also at Station 195, 1425 fathoms.

Ophiomusium validum, Ljungman. Obtained also at Station 23.

Ophiopyren longispinus, n.g., n.sp. Obtained also at Stations 23 and 33.

Ophiomitra dipsacos, n.sp. Obtained at no other locality.

Ophioscolex tropicus, n.sp. Obtained at no other locality.

ECHINOIDEA (Agassiz, Zool. pt. 9).

Dorocidaris (Cidaris) papillata (Leske). Obtained also at Stations 204, 210, 320, Canaries, and St. Paul's Rocks, 70 to 600 fathoms.

Salenia varispina, Agassiz. For distribution see Station 23.

Podocidaris sculpta, Agassiz. Obtained at no other locality by the Challenger.

ANNELIDA (M'Intosh, Zool. pt. 34).

Aphrodita intermedia, n.sp. Obtained also at Station 23.

Nothria tenuisetis, n.sp. One fragmentary specimen (either from this Station or Station 23); obtained also at Station 169.