

Rissoa, five other species undetermined.

Rissoina chesnelii (Michaud). Obtained at no other locality by the Challenger.
Recorded from West and East Indies, Mauritius, Ceylon,
and Nicobars.

„ sp. (?).

Fenella elongata, n.sp. Obtained also at Station 78, 1000 fathoms.

Actæon turritus, n.sp. One specimen; obtained at no other locality.

Ringicula peracuta, n.sp. Obtained also at Stations 56 and 122, 1075 and 350
fathoms.

Scaphander punctostriatus (Mighels). Obtained also at Stations 73 and 78, 1000
fathoms. Recorded from Arctic, North Atlantic, and Medi-
terranean.

Utriculus spatha, n.sp. Obtained at no other locality.

„ , two other species undetermined.

Cylichna ovata, Jeffreys (?). Obtained also at Stations 73, 75 (?), 78 (?), and 122,
350 to 1000 fathoms. Recorded from North-East Atlantic
("Porcupine," "Travailleur," and "Washington"). Fossil—
Middle Pliocene of Italy.

„ *discus*, n.sp. Obtained at no other locality.

„ (*Volvula*) *paupercula*, n.sp. Obtained at no other locality.

Cæcum lineicinctum, de Folin, n.sp. One specimen; obtained at no other locality.

POLYZOA (Busk, Zool. pt. 30).

Pasythea eburnea (Smitt). Obtained also at Stations 23 and 122.

Farciminaria atlantica, n.sp. Obtained also at Station 23.

BRACHIOPODA (Davidson, Zool. pt. 1).

Terebratula vitrea (Born), var. *minor*, Philippi. One specimen; obtained also at
Stations 73 and 142, 1000 and 150 fathoms. Recorded
from Arctic, North Atlantic, and Mediterranean. Fossil—
Pliocene of Sicily.

Terebratulina wyvillii, n.sp. One specimen; obtained at no other locality. Another
specimen said to be in Amsterdam Museum without name
or derivation.

Magasella incerta, n.sp. Twelve small specimens, probably immature; obtained
at no other locality.

In addition to the foregoing, the following are recorded in the Station-book:—Alcyonarian, *Archaster* sp. (?), *Myriotrochus*, several Annelids (only one reported), *Echiurus*, *Chætoderma* (?), Brachyuran.