

STATION 33.

BRACHYURA (Miers, Zool. pt. 49).

Geryon (?) *incertus*, n.sp. One injured specimen; obtained at no other locality.

LAMELLIBRANCHIATA (Smith, Zool. pt. 35).

Corbula philippii, n.sp. Obtained at no other locality by the Challenger. Recorded from Hayti.

Næra consociata, n.sp. Obtained also at Station 24.

„ *curta*, Jeffreys. Single valve; obtained also at Station 75, 450 fathoms. Recorded from North Atlantic (“Porcupine”).

„ *claviculata*, Dall. Obtained at no other locality by the Challenger. Recorded from North Atlantic (“Blake”).

„ *congenita*, n.sp. Single valve, afterwards unfortunately destroyed; obtained at no other locality.

Ervilia subcancellata, n.sp. Obtained also at Stations 113A and 120, 25 and 675 fathoms. Recorded from West Indies.

Circe bermudensis, n.sp. Obtained at no other locality.

Verticordia ornata (d'Orbigny), var. Three valves; obtained at no other locality by the Challenger. Recorded from West Atlantic, California, and China Seas.

Lucina (*Codakia*) *pecten*, Lamarck. A few small valves; obtained also at Station 113A, 25 fathoms. Recorded from North Atlantic and Mediterranean.

Cryptodon barbatus (Reeve) (?). Single valve; obtained at no other locality by the Challenger.

„ *croulinensis* (Jeffreys). Several valves; for distribution see Station VIII.

Malletia veneriformis, n.sp. Obtained also at Station 24.

Amussium dalli, n.sp. Obtained at no other locality.

„ *squamigerum*, n.sp. Obtained also at Station 24.

„ *cancellatum*, n.sp. Obtained also at Stations 24, 56, and St. Thomas.

GASTEROPODA (Watson, Zool. pt. 42).

Fissurella (*Lucapina*) *cayenensis*, Lamarck. One specimen; obtained also at Station 122, 350 fathoms. Recorded from North-West Atlantic.

Natica radiata, n.sp. Obtained also in North Atlantic, over 1000 fathoms (?).

BRACHIOPODA (Davidson, Zool. pt. 1).

Crania sp. (?). One imperfect valve.

In addition to the foregoing, the following are recorded in the Station-book:—