## AT BERMUDA. ECHINOIDEA (Agassiz, Zool. pt. 9).

Toxopneustes variegatus (Lamarck). Obtained at no other locality by the Challenger.

Spatangus purpureus, Müller. (100 fathoms); obtained also at Station 75, 50 to 90 fathoms. Recorded from East Atlantic and Caribbean Sea.

## HOLOTHURIOIDEA (Théel, Zool. pt. 39).

Synapta picta, n.sp. One specimen; obtained at no other locality.

Stichopus möbii, Semper. One specimen; obtained at no other locality by the Challenger. Recorded from West Indies.

,, haytiensis, Semper. One specimen; obtained at no other locality by the Challenger. Recorded from West Indies.

## NEMERTEA (Hubrecht, Zool. pt. 54).

Tetrastemma agricola, Willemoes-Suhm, n.sp. Numerous specimens (Mangrove Swamps, Hungry Bay); obtained at no other locality.

Cerebratulus truncatus, n.sp. (?). One specimen; obtained also at Stations 48 and 49.

## Annelida (M'Intosh, Zool. pt. 34).

Hermodice carunculata, Pallas. Littoral region; obtained also at Station 55, St. Thomas, and Cape Verdes.

Eurythoë pacifica, Kinberg (?). Several specimens (between tide-marks); obtained at no other locality by the Challenger. Recorded from coral reefs of the Pacific.

Polynoë pustulata, n.sp. Two specimens (between tide-marks); obtained at no other locality.

Nereis (Perinereis) inelanocephala, n.sp. One specimen (between tide-marks); obtained at no other locality.

Europe. Europe. Several specimens (between tide-marks); obtained also at Station 162, 38 fathoms. Recorded from Europe.

- " cirrobranchiata, n.sp. One specimen (between tide-marks); obtained at no other locality.
- " barvicensis, n.sp. One specimen (between tide-marks); obtained at no other locality.
- " sp. (?). Fragment (between tide-marks).

Aricia platycephala, n.sp. One specimen (between tide-marks); obtained at no other locality.

Cirratulus assimilis, n.sp. One specimen (between tide-marks); obtained at no other locality.