

- Sigalion buskii*, M'Intosh. One injured specimen; obtained at no other locality by the Challenger. Recorded from Shetland. STATION 75.
- Glycera tessellata*, Grube. One injured specimen; obtained also at Station 142 (?), 150 fathoms. Recorded from North Atlantic.
- „ *capitata*, CErsted. One specimen; obtained also at Station II.
- Ditrypa arietina*, Müller. Dead tubes inhabited by Gephyreans; obtained also at Station VIIIp.

## OSTRACODA (Brady, Zool. pt. 3).

- Bairdia victrix*, Brady. For distribution see Station 24.
- „ *angulata*, Brady. One or two specimens; obtained also at Stations 185 and 305, 155 and 165 fathoms. Recorded from Magellan Strait.
- Cythere dictyon*, n.sp. Widely distributed (see Station 24).
- Cytherella lata*, n.sp. A few detached valves; obtained also at Stations 24, 120, 185, and 191.

## ANOMURA (Henderson, Zool. pt. 69).

- Anapagurus pusillus*, n.sp. Several specimens (50 to 90 fathoms); obtained also at Station VIIIp. and Simon's Bay, Cape.

## BRACHYURA (Miers, Zool. pt. 49).

- Inachus leptochirus*, Leach. Three specimens (50 to 90 fathoms); obtained at no other locality by the Challenger.
- Pisa (Arctopsis) tribulus* (Linné). One specimen (50 to 90 fathoms); obtained also at Station VIIIp.
- Lambrus (Parthenolambrus) massena*, Roux. One specimen (50 to 90 fathoms); obtained also at Cape Verdes. Recorded from Mediterranean and Senegambia.
- „ ( „ ) *expansus*, Miers. Three specimens (50 to 90 fathoms); obtained at no other locality by the Challenger. Recorded from Madeira.
- Heterocrypta maltzani*, Miers. Two specimens (50 to 90 fathoms) and one specimen (450 fathoms); obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean.
- Xanthodes melanodactylus*, M.-Edwards. Numerous specimens (50 to 90 fathoms); obtained also at Canaries and Cape Verdes. Recorded from North and South Atlantic.