

- Bittium reticulatum* (Costa). Obtained also at Station VIIp.
- „ *amblyterum*, n.sp. Obtained at no other locality by the Challenger.  
Recorded from North Atlantic.
- „ *abruptum*, n.sp. Obtained at no other locality.
- „ sp. (?).
- Triforis perversa* (Linné). Obtained also at Stations VIIp. and 122.
- Cithna tenella* (Jeffreys) (?). One specimen; obtained also at Stations 78, 85, 120, and 122, 350 to 1125 fathoms. Recorded from North Atlantic and Mediterranean. Fossil—Middle Pliocene onwards.
- Rissoa fayalensis*, n.sp. Obtained at no other locality.
- „ (*Alvania*) *calathus*, Forbes and Hanley (?). One injured specimen; obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean. Fossil—Middle Pliocene.
- „ ( „ ) *cancellata* (Costa). Obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean. Fossil—Upper Miocene.
- „ ( „ ) *hispidula* (Monterosato). Obtained at no other locality by the Challenger. Recorded from Mediterranean. Fossil—Middle Pliocene onwards.
- „ ( „ ) *tarsodes*, n.sp. Obtained at no other locality.
- „ (*Setia*) *quisquiliarum*, n.sp. Obtained at no other locality.
- Actæon exilis*, Jeffreys. Obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean. Fossil—Middle Pliocene of Calabria.
- Bulla pinguicula*, Jeffreys. One broken specimen; obtained at no other locality by the Challenger. Recorded from Bay of Biscay.
- „ *semilevis*, Seguenza. Obtained also at Stations 73 and 78.
- Cylichna alba* (Brown). Obtained also at Station 122, 350 fathoms. Recorded from the whole North Atlantic and Arctic, and Japan. Fossil—Norwich Crag and Middle Pliocene of Calabria onwards.
- „ *ovata*, Jeffreys (?). Obtained also at Stations 24, 73, 78, and 122.
- Philine quadrata* (Wood) (?). (50 to 90 fathoms); obtained also at Station 78, 1000 fathoms.
- Rotella* (?) sp. (50 to 90 and 450 fathoms.)
- Minute shell, genus unknown.