

OPHIUROIDEA (Lyman, Zool. pt. 14).

STATION 85.

Ophiomitra chelys (Wyville Thomson), n.sp. Obtained also at Stations 3 and 33.
[Stated in error to be from Station 84.]

OSTRACODA (Brady, Zool. pt. 3).

Cythere dasyderma, n.sp. Widely distributed (see Station 5).

„ (?) *serratula*, n.sp. A few separate valves; obtained also at Stations 24 and 335.

Krithe producta, n.sp. Widely distributed (see Station 70).

MACRURA (Spence Bate, Zool. pt. 52).

Stochasmus exilis, n.g., n.sp. One injured specimen; obtained at no other locality.
Only species of the genus. [Stated in error to be from Station 84.]

LAMELLIBRANCHIATA (Smith, Zool. pt. 35).

Næra circinata, Jeffreys. Obtained also at Station 73.

Semele (Abra) profundorum, n.sp. Obtained also at Stations 73, 98, and 244.

Verticordia quadrata, n.sp. Single valve; obtained at no other locality.

SCAPHOPODA and GASTEROPODA (Watson, Zool. pt. 42).

Dentalium subterfissum, Jeffreys. Obtained also at Stations 78 and 120.

Siphodentalium pusillum, n.sp. Obtained at no other locality.

Cadulus gracilis, Jeffreys. Obtained also at Stations 75 and 78.

„ *tumidosus*, Jeffreys. Obtained also at Station 78.

Trochus sp. (?)

Basilissa munda, n.g., n.sp. Obtained at no other locality.

Seguenzia carinata, Jeffreys. For distribution see Station 73.

Purpura (?) sp., fry.

Pleurotoma (Mangelia) sp. (?)

Clathurella formosa (Jeffreys). Obtained also at Stations 24, 73, and 78.

„ *chariessa*, n.sp. Obtained also at Stations 24, 73, 78, and 122.

Cassis (?), two species, fry.

Natica sp. (?). Obtained also at Stations 24, 120, and 122.

Cithna tenella (Jeffreys). For distribution see Station 75.

Rissoa (Alvania) deliciosa, Jeffreys. Obtained also at Station 122, 350 fathoms.
Recorded from N.E. Atlantic and Mediterranean.

„ *amblia*, n.sp. Obtained at no other locality.

Actæon amabilis, n.sp. Obtained also at Station 73.

Bellerophina sp. (?)