

CAPE VERDES.

- Blennius sanguinolentus*, Pall. (St. Vincent); obtained at no other locality by the Challenger.
- Lepadogaster gouani*, Barnev. (St. Vincent); obtained at no other locality by the Challenger.
- Scarus chrysopterus*, Bl. (St. Vincent); obtained at no other locality by the Challenger.
- Rhomboideichthys podas*, de la Roche. (St. Vincent); obtained at no other locality by the Challenger.
- Hemirhamphus vittatus*, Val. (St. Iago); obtained at no other locality by the Challenger.
- Balistes forcipatus*, Gm. (St. Vincent); obtained at no other locality by the Challenger.
- Monacanthus setifer*, Benn. (St. Vincent); obtained also at Japan.

In the foregoing list 151 species are enumerated, of which 38 are new to science, including representatives of 7 new genera; 31 of the new species and 3 new genera were not obtained elsewhere.

ORGANISMS FROM
SURFACE-NETS.

Surface Organisms.—The following species are recorded from the surface in the neighbourhood of the Cape Verde Islands:—

OSTRACODA (Brady, Zool. pt. 3).

Halocypris brevirostris, Dana.

COPEPODA (Brady, Zool. pt. 23).

Eucalanus setiger, n.sp.

Pleuromma abdominalis (Lubbock).

Undina vulgaris, Dana.

Scolecithrix danæ (Lubbock).

Euchæta prestandreae, Philippi.

Candace pectinata, Brady.

„ *pachydactyla*, Dana.

Temora dubia (Lubbock).

Pontella detruncata (Dana).

„ *acutifrons* (Dana).

„ *plumata* (Dana).

Corycaeus pellucidus, Dana.

Oncae obtusa (Dana).

Saphirina ovalis, Dana.

„ *serrata*, n.sp.

AMPHIPODA (Stebbing, Zool. pt. 67).

Synopia schéeleana, Bovallius.

Gammaropsis atlantica, n.sp.

Dairella bovallii, n.sp.

Phronima megalodous, n.sp.

Hyperia schizogeneios, n.sp.

Brachyscelus mediterranea (Claus).

Lycæa vincentii, n.sp.

STOMATOPODA (Brooks, Zool. pt. 45).

Alima gracilis, M.-Edwards [= larva of *Squilla*].

MACRURA (Spence Bate, Zool. pt. 52).

Phyllosoma furcicaudatum, n.sp.

„ *verdense*, n.sp.

Sergestes edwardsii, Krøyer.

„ *dissimilis*, n.sp.

Diaphoropus longidorsalis, n.g., n.sp.