

BAHIA. CRINOIDEA (Carpenter, Zool. pt. 60).

- Antedon dübeni*, Böhlische. One specimen (20 fathoms); obtained at no other locality by the Challenger. Recorded from Rio Janeiro and Abrolhos (?).
- „ *carinata* (Lamarck). Several specimens (7 to 20 fathoms); obtained at no other locality by the Challenger. A widely-distributed species.
- Actinometra meridionalis* (Agassiz). Fourteen specimens (7 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from West Atlantic.
- „ *lineata*, n.sp. Eight specimens (7 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from West Atlantic.

ASTEROIDEA (Sladen, Zool. pt. 51).

- Astropecten brasiliensis*, Müller and Troschel. Several specimens (7 to 20 fathoms); obtained also at Fernando Noronha.
- Luidia alternata* (Say), Lütken. (7 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from West Indies.
- „ *clathrata* (Say), Lütken. (7 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from West Atlantic.
- Narcissia trigonaria*, n.sp. Obtained at no other locality.
- Echinaster spinosus* (Retzius), Müller and Troschel. (7 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from West Atlantic.

OPHIUROIDEA (Lyman, Zool. pt. 14).

- Ophiura brevispina*, Say. Obtained at no other locality by the Challenger.
- „ *appressa*, Say. Obtained at no other locality by the Challenger.
- „ *cinerea* (Müller and Troschel). (7 to 20 fathoms); obtained also at St. Thomas.
- Ophiozona impressa* (Lütken). (7 to 20 fathoms); obtained at no other locality by the Challenger.
- Ophiactis mülleri*, Lütken, var. *quinqueradia*, Lyman. (7 to 20 fathoms); obtained also at Station 122.
- Ophionereis reticulata* (Say). (7 to 20 fathoms); obtained also at Bermuda.
- Ophiothrix angulata* (Say), var. (7 to 20 fathoms); obtained also at Station 36 and Fernando Noronha.