

TRISTAN DA
CUNHA.

Tentorium semisuberites (Schmidt). Four specimens of dwarf variety (60 to 90 fathoms); obtained also at Stations 49 and 50.

Latrunculia (?) *acerata*, n.sp. One injured specimen (60 fathoms); obtained at no other locality.

TETRACTINELLIDA (Sollas, Zool. pt. 63).

Pachastrella abyssi, Schmidt. Two specimens (110 fathoms); obtained at no other locality by the Challenger. Recorded from North Atlantic ("Porcupine") and Gulf of Mexico.

CALCAREA (Poléjaeff, Zool. pt. 24).

Sycon raphanus, Schmidt, var. *tergestinum*, Haeckel. One specimen (60 to 90 fathoms); obtained also at Station 209, 95 fathoms. Recorded from Adriatic.

Pericharax carteri, n.g., n.sp. One specimen and fragment (60 fathoms); obtained at no other locality. Only species of the genus.

ALCYONARIA (Wright and Studer, Zool. pts. 64 and 81).

Clavularia cylindrica, n.sp. One specimen (100 to 150 fathoms); obtained at no other locality.

Sympodium glomeratum, n.sp. One specimen (100 to 150 fathoms); obtained at no other locality.

Sarakka crassa, Danielssen. One specimen (100 to 550 fathoms); obtained at no other locality by the Challenger. Recorded from North Atlantic.

Sarcophytum atlanticum, n.sp. One specimen (60 fathoms); obtained at no other locality.

Thouarella affinis, n.sp. One specimen (55 to 70 fathoms); obtained at no other locality.

Amphilaphis regularis, n.g., n.sp. Two specimens (75 to 150 fathoms); obtained at no other locality. Only species of the genus.

Clematissa verrilli, n.g., n.sp. (360 fathoms); obtained at no other locality.

ACTINIARIA (Hertwig, Zool. pts. 15 and 73).

Tealia bunodiformis, n.sp. Three specimens (shore); obtained at no other locality.

Palythoa anguicomis, Norman. (60 to 90 fathoms); obtained at no other locality by the Challenger.

„ sp. (?). (60 to 150 fathoms).