

TRISTAN DA
CUNHA.

ECHINOIDEA (Agassiz, Zool. pt. 9).

Arbacia dufresnii (Bl.). (100 to 150 fathoms); obtained also at Stations 304 and 308, 45 and 175 fathoms.

Echinus elegans (Düben and Koren). (1100 fathoms); obtained also at Stations 46 and 219.

OSTRACODA (Brady, Zool. pt. 3).

Bairdia villosa, n.sp. (100 to 150 fathoms); obtained also at Stations 145, 149, and 162, 38 to 150 fathoms.

Cythere impluta, n.sp. (100 to 150 fathoms); obtained also at Station 316, 6 fathoms.

Cytherella punctata, Brady. (100 to 150 fathoms); obtained also at Stations 167, 191, 305, and Port Jackson, 2 to 580 fathoms. Recorded from the Levant (?).

CIRRIPIEDIA (Hoek, Zool. pt. 25).

Scalpellum carinatum, n.sp. Two specimens (1000 fathoms); obtained at no other locality.

„ *africanum*, n.sp. Four specimens (100 fathoms); obtained at no other locality.

„ *elongatum*, n.sp. (Depth not given); obtained also at Stations 164B and 169, 410 and 700 fathoms.

„ *eximium*, n.sp. One specimen attached to pumice-stone (1000 fathoms); obtained at no other locality.

AMPHIPODA (Stebbing, Zool. pt. 67).

Metopa crenatipalmata, n.sp. One specimen (100 to 150 fathoms); obtained also at Station 313, 55 fathoms.

Podocerus tristanensis, n.sp. Two specimens (110 fathoms); obtained at no other locality.

Caprellinoides tristanensis, n.g., n.sp. One specimen (110 fathoms); obtained at no other locality. Only species of the genus.

Æginella tristanensis, n.sp. One specimen (110 fathoms); obtained at no other locality.

ISOPODA (Beddard, Zool. pt. 48).

Janira tristani, n.sp. Two specimens (100 to 150 fathoms); obtained at no other locality.