

TRISTAN DA
CUNHA.

GASTEROPODA (Watson, Zool. pt. 42).

Emarginula sp. (?).

Scissurella aëdonia, n.sp. (100 to 150 fathoms); obtained also at Station 122.

„, two other species undetermined.

Scalaria philtata, n.sp. (100 to 150 fathoms); obtained at no other locality.

Murex (Pseudomurex) aëdonius, n.sp. (100 to 150 fathoms); obtained at no other locality.

Buccinum (?) sp. (100 to 150 fathoms).

Pleurotoma (Thesbia) eritima, n.sp. (100 to 150 fathoms); obtained at no other locality.

Triton (Simpulum) philomelæ, n.sp. (100 to 150 fathoms); obtained at no other locality.

Ranella (Argobuccinum) argus (Gmelin). (Shore and 100 to 150 fathoms); obtained at no other locality by the Challenger. A widely-distributed species.

Crepidula onyx, Sowerby. One specimen (shore); obtained at no other locality by the Challenger. Recorded from Panama and Mazatlan.

Odostomia (Turbonilla) philomelæ, n.sp. (100 to 150 fathoms); obtained at no other locality.

Bittium pigrum, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ *lusciniæ*, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ *philomelæ*, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ *delicatum*, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ *aëdonium*, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ sp. (?). (100 to 150 fathoms).

Triforis hebes, n.sp. Three specimens (100 to 150 fathoms); obtained at no other locality.

Rissoa (Alvania) lusciniæ, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ (*Onoba*) *arenaria* (Mighels). One specimen (100 to 150 fathoms); obtained at no other locality by the Challenger. Recorded from Arctic and North Atlantic.

„ („) *aëdonis*, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ (*Ceratia*) *glaphyra*, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ („) *macra*, n.sp. (100 to 150 fathoms); obtained at no other locality.

„ (*Setia*) *philomelæ*, n.sp. (100 to 150 fathoms); obtained at no other locality.