

CAPE OF GOOD  
HOPE.

## OPHIUROIDEA (Lyman, Zool. pt. 14).

*Ophiura tongana* (Lütken) (?). (Simon's Bay, 10 to 20 fathoms); obtained at no other locality by the Challenger.

*Ophiactis carnea*, Ljungman. (Simon's Bay, 10 to 20 fathoms); obtained at no other locality by the Challenger.

*Amphiura incana*, n.sp. (Simon's Bay, 10 to 20 fathoms); obtained at no other locality.

*Ophiocoma scolopendrina* (Lamarek). (Simon's Bay, 10 to 20 fathoms); obtained also at Tongatabu, Fiji, and Philippines.

*Ophiothrix triglochis*, Müller and Troschel. (Simon's Bay, 5 to 18 fathoms); obtained at no other locality by the Challenger.

*Gorgonocephalus verrucosus* (Lamarek). (Simon's Bay, 10 to 20 fathoms); obtained at no other locality by the Challenger.

## ECHINOIDEA (Agassiz, Zool. pt. 9).

*Echinus angulosus* (Leske). (Simon's Bay, 10 to 20 fathoms); obtained at no other locality by the Challenger.

*Lovenia elongata*, Gray. (Simon's Bay); obtained also at Stations 188 and 212, 10 to 28 fathoms. Recorded from Gulf of California and Philippines.

*Brissopsis lyrifera* (Forbes). (Simon's Bay, 5 to 18 fathoms); obtained also at Stations 141 and 142, 98 and 150 fathoms. A widely-distributed species.

## HOLOTHURIOIDEA (Théel, Zool. pt. 39).

*Cucumaria discolor*, n.sp. One specimen (Simon's Bay, 10 to 20 fathoms); obtained at no other locality.

„ *insolens*, n.sp. Numerous specimens (Simon's Bay, 10 to 20 fathoms); obtained at no other locality.

*Holothuria africana*, n.sp. One specimen (Simon's Bay, 10 to 20 fathoms); obtained at no other locality.

## GEPHYREA (Selenka, Zool. pt. 36).

*Phascolosoma capense*, Teuscher. Numerous specimens (Sea Point, shallow water); obtained at no other locality by the Challenger. Recorded from the Cape.

## ANNELIDA (M'Intosh, Zool. pt. 34).

*Euphrosyne capensis*, Kinberg. Many specimens (Sea Point, under stones between tide-marks); obtained at no other locality by the Challenger. Recorded from the Cape and St. Paul.