

CAPE OF GOOD
HOPE.

- Bairdia ovata*, Bosquet (?). (Simon's Bay, 15 to 20 fathoms); obtained also at Station 167, 150 fathoms.
- Cythere exilis*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.
- „ *flabellucostata*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.
- „ *craticula*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.
- „ *stolonifera*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.
- „ *lepralioides*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained also at Station 142, 150 fathoms.
- Loxoconcha subrhomboidea*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.
- Xestoleberis africana*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.
- Cytherura mucronata*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.
- „ *clausi*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.
- Cytherella dromedaria*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.

AMPHIPODA (Stebbing, Zool. pt. 67).

- Lysianax variegatus* (Stimpson). Several specimens (Simon's Bay, 18 fathoms); obtained at no other locality by the Challenger. Recorded from the Cape.

ISOPODA.

Fourteen specimens, belonging to five species undetermined (Simon's Bay and Cape Town).

ANOMURA (Henderson, Zool. pt. 69).

- Dromidia spongiosa*, Stimpson. One specimen (Simon's Bay, 10 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from the Cape and St. Paul.
- Pseudodromia latens*, Stimpson. One specimen (Simon's Bay, 10 to 12 fathoms); obtained at no other locality by the Challenger. Recorded from the Cape.
- Diogenes brevirostris*, Stimpson. One specimen (Simon's Bay, 10 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from the Cape.