

Fusus verruculatus, Lamarck. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from East Indies (?).

Cominella porcata (Gmelin). (Simon's Bay, 10 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from the Cape.

Ancilla (Anaulax) obtusa (Swainson). (Simon's Bay, 15 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from the Cape and East Africa.

Columbella mercatoria (Linné). (Sea Point); obtained also at Fernando Noronha.

Mitra sp. (?). (Sea Point).

Marginella (Glabella) chrysea, n.sp. (Sea Point); obtained at no other locality.

Strombus lentiginosus, Linné. (Simon's Bay); obtained at no other locality by the Challenger. Recorded from East Africa throughout Indian Ocean to Australia.

Turritella knysnaënsis, Krauss. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from the Cape.

Odostomia sp. (?). (Simon's Bay, 15 to 20 fathoms).

Cerithium vulgatum, Bruguière. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from Black Sea, Mediterranean, and North Atlantic. Fossil—Middle Miocene of Italy onwards.

Rissoa (Onoba) fenestrata, Krauss. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from the Cape.

Aplysia depilans, Linné. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean.

Dolabrifera triangularis, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained at no other locality.

Coralliophila wahlbergi (Krauss). (Sea Point, shallow water). [See Smith, *Proc. Zool. Soc. Lond.*, 1891, pp. 436, 437].

POLYPLACOPHORA (Haddon, Zool. pt. 43).

Ischnochiton viridulus (Couthouy). One specimen (Cape Town); obtained at no other locality by the Challenger. Recorded from the Cape.

Chiton nigrovirescens, Blainville. Six specimens (Sea Point, shore); obtained at no other locality by the Challenger. Recorded from the Cape.