and sent it down again. Three hauls were taken with the dredge, and the swabs were crowded with specimens of Brissopsis lyrifera. At 5.45 P.M. made all plain sail. Weather fine, almost a calm; sea smooth.

STATION 141.

The following species are recorded in the Zoological Reports from the dredge at this Station :-

ANIMALS FROM DREDGE.

#### OPHIUROIDEA (Lyman, Zool. pt. 14).

Ophioglypha costata, n.sp. Obtained also at Station 142, 150 fathoms.

Amphiura capensis, Ljungman. Obtained at no other locality by the Challenger.

- dilatata, n.sp. Obtained at no other locality.
- squamata (Delle Chiaje), Sars. Obtained also at Station 163, 120 fathoms. A widely-distributed species.

#### ECHINOIDEA (Agassiz, Zool. pt. 9).

Spatangus raschi, Lovén. Obtained also at Station 142, 150 fathoms. Recorded from East Atlantic.

Brissopsis lyrifera (Forbes). Many specimens; obtained also at Simon's Bay, Cape, and Station 142.

## HOLOTHURIOIDEA (Théel, Zool. pt. 39).

Cucumaria capensis, n.sp. One specimen; obtained also at Station 142, 150 fathoms.

## Annelida (M'Intosh, Zool. pt. 34).

Polynoë capensis, n.sp. Two specimens; obtained at no other locality.

Syllis capensis, n.sp. One small specimen; obtained at no other locality.

Lumbriconereis pettigrewi, n.sp. Several specimens; obtained at no other locality.

Scalibregma inflatum, Rathke, var. (?). Numerous specimens; obtained also at Station 169, 700 fathoms.

One fragmentary specimen; obtained at no other Ranzania (?) capensis, n.sp. locality.

Prionospio capensis, n.sp. One fragmentary specimen; obtained at no other locality.

Two fragmentary specimens; obtained at no other Praxilla capensis, n.sp. locality.

# Schizopoda (Sars, Zool. pt. 37).

Lophogaster typicus, Sars. One or two specimens; obtained also at Station 142, 150 fathoms. Recorded from North Atlantic.