

- Psolus ephippifer*, Wyville Thomson, n.sp. Numerous specimens (depth not given), and one specimen (310 fathoms); obtained also at Stations 149, 150, and 151, 20 to 150 fathoms. STATIONS 144A TO 145A.
- Pseudostichopus mollis*, n.g., n.sp. Two specimens (50 to 75 fathoms); obtained also at Stations 309 and 311, 140 and 245 fathoms.

## NEMERTEA (Hubrecht, Zool. pt. 54).

- Amphiporus marioni*, n.sp. One specimen (69 fathoms); obtained also at Station 149, 120 fathoms.
- Cerebratulus longifissus*, n.sp. Two specimens (69 fathoms); obtained at no other locality.

## ANNELIDA (M'Intosh, Zool. pt. 34).

- Lætmonice producta*, Grube, var. *wyvillei*, nov. Several specimens (50 to 150 fathoms); obtained also at Station 157, 1950 fathoms.
- Polyeunoa lævis*, n.g., n.sp. Numerous specimens (310 fathoms); obtained also at Station 310, 400 fathoms. Only species of the genus.
- Lagisca antarctica*, n.sp. Numerous specimens (69 fathoms); obtained also at Station 149, 127 fathoms.
- „ *magellanica*, n.sp., var. *grubei*, nov. Several specimens (310 fathoms); the species obtained also at Stations 149, 308, and 310, 127 to 400 fathoms.
- Exogone heterosetosa*, n.sp. One specimen (69 fathoms); obtained at no other locality.
- Nereis (Platynereis) catoni*, M'Intosh. One fragmentary specimen (69 fathoms); obtained also at Station 149, Fernando Noronha, and Falkland Islands.
- Eunice ærstedii*, Stimpson (?). Several specimens (69 fathoms); obtained also at Stations 45 and 49.
- „ *edwardsi*, n.sp. One specimen (140 fathoms); obtained at no other locality.
- Terebella (Lanice) flabellum*, Baird. (69 and 150 fathoms); obtained also at Station 163A, 150 fathoms. Recorded from Narçon Island.
- Neottis antarctica*, M'Intosh. (69 fathoms); obtained also at Stations 149, 150, 151, and 313, 20 to 150 fathoms.
- Serpula narconensis*, Baird. Numerous specimens (69 fathoms); obtained also at Stations 149, 151, and 308, 60 to 175 fathoms. Recorded from Narçon Island, Kerguelen, and Magellan Strait.